

Twisted Particle Filters

Nick Whiteley* and Anthony Lee†

October 2, 2012

Abstract

We investigate alternative sampling laws for particle algorithms and the influence of these laws on the efficiency of particle approximations of marginal likelihoods in hidden Markov models. Amongst a broad class of candidates we characterize the essentially unique family of particle system transition kernels which is optimal with respect to an asymptotic-in-time variance growth rate criterion. The sampling structure of the algorithm defined by these optimal transitions turns out to be only quite subtly different from standard algorithms and yet the fluctuation properties of the estimates it provides are, in some ways, dramatically different. The structure of the optimal transition suggests a new class of algorithms, which we term "twisted" particle filters, and which we validate with asymptotic analysis of a more traditional nature, in the regime where the number of particles tends to infinity.

1 Introduction

A hidden Markov model (HMM) is a bi-variate process $\{(X_n, Y_n); n \geq 0\}$ where the signal process $(X_n; n \geq 0)$ is a Markov chain and each observation Y_n is conditionally independent of the rest of the bi-variate process given X_n . Each X_n is valued in a state-space $\mathbf{X}$ endowed with a σ -algebra $\mathcal{X}$, and each Y_n is valued in an observation space $\mathbf{Y}$ endowed with a σ -algebra $\mathcal{Y}$. This paper is centrally motivated by the task of computing the marginal likelihood of an observation sequence $(Y_0, Y_1, \dots)$, under some assumed probability model for the joint process $\{(X_n, Y_n); n \geq 0\}$; let μ_0 and f be respectively a probability distribution and a Markov kernel on $(\mathbf{X}, \mathcal{X})$, and let g be a Markov kernel acting from $(\mathbf{X}, \mathcal{X})$ to $(\mathbf{Y}, \mathcal{Y})$, with $g(x, \cdot)$ admitting a strictly positive density, similarly denoted by $g(x, y)$, with respect to some dominating σ -finite measure. The hidden Markov model specified by μ_0 , f and g , is

$$\begin{aligned} X_0 &\sim \mu_0(\cdot), \quad Y_0 | \{X_0 = x_0\} \sim g(x_0, \cdot), \\ X_n | \{X_{n-1} = x_{n-1}\} &\sim f(x_{n-1}, \cdot), \quad Y_n | \{X_n = x_n\} \sim g(x_n, \cdot), \quad n \geq 1. \end{aligned} \quad (1)$$

In practice one is presented with $(Y_0, Y_1, \dots)$ but typically does not know which particular μ_0, f and g specify the true law of this process. More broadly, one may not be sure that $(Y_0, Y_1, \dots)$ are observations from an HMM at all, but seeks to compare and fit different models to these data and it is here that the computation of the marginal likelihood is especially desirable.

Let $\Omega := \mathbf{Y}^{\mathbb{Z}}$ be the set of doubly infinite sequences valued in $\mathbf{Y}$. We shall write $Y_n(\omega) = \omega(n)$ for $\omega = \{\omega(n)\}_{n \in \mathbb{Z}} \in \Omega$. For any $\omega \in \Omega$ we may take as a recursive definition of the (one-step-ahead) *prediction filters*, the sequence of distributions $(\pi_n^\omega; n \geq 0)$ following

$$\begin{aligned} \pi_0^\omega &:= \mu_0, \\ \pi_n^\omega(A) &:= \frac{\int_{\mathbf{X}} \pi_{n-1}^\omega(dx) g(x, Y_{n-1}(\omega)) f(x_{n-1}, A)}{\int_{\mathbf{X}} \pi_{n-1}^\omega(dx) g(x, Y_{n-1}(\omega))}, \quad A \in \mathcal{X}, \quad n \geq 1. \end{aligned} \quad (2)$$

We are centrally concerned with the sequence $(Z_n^\omega; n \geq 0)$ defined by

$$Z_0^\omega := 1, \quad Z_n^\omega := Z_{n-1}^\omega \int_{\mathbf{X}} \pi_{n-1}^\omega(dx) g(x, Y_{n-1}(\omega)), \quad n \geq 1, \quad (3)$$

*University of Bristol, School of Mathematics, University Walk, Bristol, BS8 1TW, UK

†University of Warwick, Department of Statistics, Coventry, CV4 7AL, UK

which can, in principle, be computed sequentially as a by-product of the filtering recursion (2). Under the model (1), π_n^ω is interpreted as the conditional distribution of X_n given $Y_{0:n-1}(\omega)$; and Z_n^ω is the marginal likelihood evaluated at the point $Y_{0:n-1}(\omega)$. If we write $\mathbf{E}$ for expectation with respect to the law of the Markov chain (X_n) under the assumed model (1), i.e. $X_n|X_{n-1} \sim f(X_{n-1}, \cdot)$ and $X_0 \sim \mu_0$, then we have the standard identity

$$Z_n^\omega = \mathbf{E} \left[\prod_{p=0}^{n-1} g(X_p, Y_p(\omega)) \right].$$

The simplest particle filter, known as the “bootstrap” algorithm [Gordon et al., 1993], is given below.

Algorithm 1 Bootstrap particle filter

For $n = 0$,

Sample $(\zeta_0^i)_{i=1}^N \stackrel{\text{iid}}{\sim} \mu_0$,

Report $Z_0^{\omega, N} = 1$.

For $n \geq 1$,

Report $Z_n^{\omega, N} = Z_{n-1}^{\omega, N} \cdot \frac{1}{N} \sum_{j=1}^N g(\zeta_{n-1}^j, Y_{n-1}(\omega))$,

Sample $(\zeta_n^i)_{i=1}^N \mid (\zeta_{n-1}^i)_{i=1}^N \stackrel{\text{iid}}{\sim} \frac{\sum_{j=1}^N g(\zeta_{n-1}^j, Y_{n-1}(\omega)) f(\zeta_{n-1}^j, \cdot)}{\sum_{j=1}^N g(\zeta_{n-1}^j, Y_{n-1}(\omega))}$,

There are now many theoretical analyses of particle filtering algorithms which validate their application. Consistency and fluctuation properties of particle algorithms in the regime $N \rightarrow \infty$ are well understood [Del Moral and Guionnet, 1999, Chopin, 2004, Künsch, 2005, Douc and Moulines, 2008]; stability properties of particle approximations have been expressed through finite- N error bounds [Del Moral and Guionnet, 2001, Cérou et al., 2011, Whiteley, 2011], time-uniform convergence [van Handel, 2009] and control on $N \rightarrow \infty$ asymptotic variance expressions [Del Moral and Jacod, 2001, Favetto, 2012, Whiteley, 2011, Douc et al., 2012].

In the present work we conduct a study of $Z_n^{\omega, N}$, and related quantities, in the regime where N is fixed and $n \rightarrow \infty$. Our overall aim is not just to validate algorithms, but to rigorously address more comparative questions of how and why one algorithm may out-perform another, and how it may be possible to modify standard algorithms in order to improve performance. Our study is formulated in a generic framework which accommodates various standard particle algorithms and novel extensions. Much of the structure we uncover is non-standard, and the algorithms we devise are new, so for the sake of an accessible introduction we shall now summarize some of our intentions and findings in the context of the bootstrap particle filter as per Algorithm 1 (more precise and complete statements will be given in a general setting later).

Writing $\mathbf{E}_N^\omega$ for expectation with respect to the law of the bootstrap particle filter processing a particular fixed observation sequence $\omega \in \Omega$, the well known lack-of-bias property [Del Moral, 2004, Chapter 9] reads:

$$\mathbf{E}_N^\omega [Z_n^{\omega, N}] = Z_n^\omega, \quad (4)$$

and holds for any $n \geq 0$ and $N \geq 1$. The main theme of this paper is exploration and analysis of novel particle algorithms arising through changes of measure on the left hand side of (4); we shall consider alternative sampling laws which govern the transitions of the particle system as a whole and which generalize the sampling recipe in Algorithm 1. The resulting approximations of Z_n^ω will be of the form

$$\tilde{Z}_n^{\omega, N} := Z_n^{\omega, N} \cdot \prod_{p=1}^n \phi_p^{\omega, N}, \quad n \geq 1 \quad (5)$$

where $Z_n^{\omega, N}$ is exactly the same functional of the particles as in Algorithm 1 and $\{\phi_n^{\omega, N}; n \geq 1\}$ is a sequence of functionals chosen such that, if we write $\tilde{\mathbf{E}}_N^\omega$ for expectation under the (as yet unspecified) alternative sampling law, then the lack of bias property is preserved:

$$\tilde{\mathbf{E}}_N^\omega [\tilde{Z}_n^{\omega, N}] = Z_n^\omega. \quad (6)$$

Our main objective is to identify “good” choices of alternative sampling laws, possibly allowing the transitions of the particles to depend on past and/or future observations. Our criterion for performance will involve certain fluctuation properties of $\tilde{Z}_n^{\omega,N}$; we shall obtain a mathematically fruitful study of the normalized second moment of $\tilde{Z}_n^{\omega,N}$, in the regime where N is fixed and $n \rightarrow \infty$, in an ω -pathwise fashion. This is where the explicit instantiation of Ω will be utilized, and certain ergodicity hypotheses will be introduced to facilitate our $n \rightarrow \infty$ limit theory.

For now, let us still consider $\omega \in \Omega$ as fixed. Then under the probability law corresponding to Algorithm 1, the generations of the particle system, $\zeta_0, \zeta_1, \dots$ with $\zeta_n := (\zeta_n^1, \dots, \zeta_n^N)$ form an $\mathbf{X}^N$ -valued time-inhomogeneous Markov chain. Let $\{\mathbf{M}^\omega; \omega \in \Omega\}$ be the family of Markov kernels such that for each $\omega \in \Omega$, $\mathbf{M}^\omega : \mathbf{X}^N \times \mathcal{X}^{\otimes N} \rightarrow [0, 1]$ is given by

$$\mathbf{M}^\omega(x, dz) = \prod_{i=1}^N \frac{\sum_{j=1}^N g(x^j, Y_0(\omega)) f(x^j, dz^i)}{\sum_{j=1}^N g(x^j, Y_0(\omega))}, \quad (7)$$

with $x = (x^1, \dots, x^N) \in \mathbf{X}^N$ and $z = (z^1, \dots, z^N) \in \mathbf{X}^N$. Let $\theta : \Omega \rightarrow \Omega$ be the shift operator, $(\theta\omega)(n) := \omega(n+1)$, $n \in \mathbb{Z}$, $\omega \in \Omega$, so that for example $Y_0(\theta\omega) = Y_1(\omega)$. The n -fold iterate of θ will be written θ^n with $\theta^0 = Id$. It is then clear that the sampling steps of Algorithm 1 implement:

$$\zeta_0 \sim \mu_0^{\otimes N}, \quad \zeta_n | \zeta_{n-1} \sim \mathbf{M}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot), \quad n \geq 1. \quad (8)$$

Variance Growth Rates

We shall study algorithms which involve sampling the particle system under alternatives to (8) and which in general may be of a very different nature to (7): for a family of Markov kernels $\{\tilde{\mathbf{M}}^\omega; \omega \in \Omega\}$ belonging to a broad class of candidates and which may depend on ω in a rather general fashion, but subject to

$$\mathbf{M}^\omega(x, \cdot) \ll \tilde{\mathbf{M}}^\omega(x, \cdot),$$

and other regularity conditions, we shall consider sampling the particle system according to

$$\zeta_0 \sim \mu_0^{\otimes N}, \quad \zeta_n | \zeta_{n-1} \sim \tilde{\mathbf{M}}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot), \quad n \geq 1, \quad (9)$$

and simply setting

$$\phi_n^{\omega,N} := \frac{d\mathbf{M}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot)}{d\tilde{\mathbf{M}}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot)}(\zeta_n), \quad n \geq 1.$$

Then letting $\tilde{\mathbb{E}}_N^\omega$ denote expectation under the Markov law (9), and with $\tilde{Z}_n^{\omega,N}$ as in (5), we of course achieve (6).

Let Ω be endowed with the product σ -algebra $\mathcal{F} = \mathcal{Y}^{\otimes \mathbb{Z}}$ and let $\mathbb{P}$ be a probability measure on $(\Omega, \mathcal{F})$. The requirement to compute Z_n^ω typically arises when calibrating and comparing HMMs, so it is very important that our study accommodates model mis-specification and we stress that $\mathbb{P}$ is not necessarily the measure on observation sequences induced by the particular HMM (1), nor necessarily the measure induced by any HMM. Under the assumptions that θ is $\mathbb{P}$ -preserving and ergodic, and under certain other regularity conditions, application of our first main result, Proposition 4, establishes, for any *fixed* $N \geq 1$, existence of a deterministic constant $\Upsilon_N(\tilde{\mathbf{M}})$, depending on $\tilde{\mathbf{M}} = \{\tilde{\mathbf{M}}^\omega; \omega \in \Omega\}$ such that

$$\frac{1}{n} \log \frac{\tilde{\mathbb{E}}_N^\omega \left[\left(\tilde{Z}_n^{\omega,N} \right)^2 \right]}{(Z_n^\omega)^2} \rightarrow \Upsilon_N(\tilde{\mathbf{M}}), \quad \text{as } n \rightarrow \infty, \quad \text{for } \mathbb{P} - a.a. \omega. \quad (10)$$

When $\Upsilon_N(\tilde{\mathbf{M}})$ exists, it must be the case that $\Upsilon_N(\tilde{\mathbf{M}}) \geq 0$, because variance is non-negative and the lack of bias property, (6), holds. We shall see that it is typically the case that $\Upsilon_N(\tilde{\mathbf{M}}) > 0$.

Optimal Sampling

Our second main result (Theorem 1) identifies, for any fixed $N \geq 1$ and amongst the class of candidates, the essentially unique choice of the family $\{\widetilde{\mathbf{M}}^\omega; \omega \in \Omega\}$ which achieves $\Upsilon_N(\widetilde{\mathbf{M}}) = 0$. It turns out that this optimal choice of $\{\widetilde{\mathbf{M}}^\omega; \omega \in \Omega\}$ arises from a particular form of re-weighting applied to each transition kernel $\mathbf{M}^\omega$, and is defined in terms of a family of functions $\{h^\omega : \mathbf{X} \rightarrow \mathbb{R}_+; \omega \in \Omega\}$ which are, in abstract terms, generalized eigen-functions associated with algebraic structures underlying the particle algorithm. These functions have a particular probabilistic interpretation in the context of HMMs, which we shall discuss in section 4. Theorem 1 establishes that for any $N \geq 1$, $\Upsilon_N(\widetilde{\mathbf{M}}) = 0$ if and only if, for $\mathbb{P}$ -almost all $\omega \in \Omega$ there exists a set $A_\omega \in \mathcal{X}^{\otimes N}$ such that A_ω^c is null (with respect to an as yet un-named measure) and for any $x \in A_\omega$,

$$\widetilde{\mathbf{M}}^\omega(x, B) = \frac{\int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}, \quad \text{for all } B \in \mathcal{X}^{\otimes N}, \quad (11)$$

where

$$\mathbf{h}^\omega : x = (x^1, \dots, x^N) \in \mathbf{X}^N \mapsto \frac{1}{N} \sum_{i=1}^N h^\omega(x^i) \in \mathbb{R}_+.$$

Having begun this introductory discussion by stating the algorithmic definition of the bootstrap particle filter, then writing the more probabilistic representation (7)-(8), we shall now reverse this programme and proceed to write down the algorithm which arises when sampling under a re-weighted Markov transition such as (11). In the rare-event and large deviations literatures, the action of re-weighting Markov kernels using non-negative eigen-functions is generically referred to as “twisting”. Since in the present context we are applying re-weighting to the transitions of the entire particle system, we shall adopt this terminology and consider a class of algorithms which we refer to as *twisted particle filters*.

Twisted Particle Filters

The form of the optimal transition (11), where $\mathbf{M}^\omega$ is re-weighted by an additive, non-negative functional, leads us to consider a new class of particle algorithms. Consider a family of functions $\{\psi^\omega : \mathbf{X} \rightarrow \mathbb{R}_+; \omega \in \Omega\}$ and let $\{\widetilde{\mathbf{M}}^\omega; \omega \in \Omega\}$ be defined by

$$\widetilde{\mathbf{M}}^\omega(x, dx') \propto \mathbf{M}^\omega(x, dx') \psi^{\theta\omega}(x'), \quad (12)$$

with

$$\psi^\omega : x = (x^1, \dots, x^N) \in \mathbf{X}^N \mapsto \frac{1}{N} \sum_{i=1}^N \psi^\omega(x^i) \in \mathbb{R}_+. \quad (13)$$

This setup clearly admits the optimal transition ($\psi^\omega = h^\omega$) and the standard transition (take $\psi^\omega = c$, for some positive constant c), as special cases. Then introducing

$$\widetilde{g}^\omega(x) := g(x, Y_0(\omega)) \int_{\mathbf{X}} f(x, dz) \psi^{\theta\omega}(z),$$

we observe that

$$\phi_n^{\omega, N} = \frac{d\mathbf{M}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot)}{d\widetilde{\mathbf{M}}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot)}(\zeta_n) = \left[\frac{1}{N^{-1} \sum_{i=1}^N g(\zeta_{n-1}^i, Y_{n-1}(\omega))} \right] \frac{\sum_{i=1}^N \widetilde{g}^{\theta^{n-1}\omega}(\zeta_{n-1}^i)}{\sum_{i=1}^N \psi^{\theta^n\omega}(\zeta_n^i)}.$$

Since ψ^ω is an additive functional, it is clear that $\widetilde{\mathbf{M}}^\omega$ as per (12) will be of mixture form, and introducing the ω -dependent Markov kernel

$$\widetilde{f}^\omega(x, dx') := \frac{f(x, dx') \psi^{\theta\omega}(x')}{\int_{\mathbf{X}} f(x, dz) \psi^{\theta\omega}(z)},$$

the procedure of sampling from (9) and evaluating $\tilde{Z}_n^{\omega,N}$ can be implemented through Algorithm 2 (here K_n and A_n are just some auxiliary random variables employed for algorithmic purposes).

Algorithm 2 Twisted bootstrap particle filter

For $n = 0$,

Sample $(\zeta_0^i)_{i=1}^N \stackrel{\text{iid}}{\sim} \mu_0$,

Report $\hat{Z}_0^{\omega,N} = 1$.

For $n \geq 1$,

Sample K_n from the uniform distribution on $\{1, \dots, N\}$,

Sample A_n from the distribution on $\{1, \dots, N\}$ with probabilities proportional to

$$\left\{ \tilde{g}^{\theta^{n-1}\omega}(\zeta_{n-1}^1), \dots, \tilde{g}^{\theta^{n-1}\omega}(\zeta_{n-1}^N) \right\}$$

Sample $\zeta_n^{K_n} \mid \left\{ A_n, K_n, (\zeta_{n-1}^i)_{i=1}^N \right\} \sim \tilde{f}^{\theta^n \omega}(\zeta_{n-1}, \cdot)$

Sample $(\zeta_n^i)_{i \neq K_n} \mid \left\{ K_n, (\zeta_{n-1}^i)_{i=1}^N \right\} \stackrel{\text{iid}}{\sim} \frac{\sum_{j=1}^N g(\zeta_{n-1}^j, Y_{n-1}(\omega)) f(\zeta_{n-1}^j, \cdot)}{\sum_{j=1}^N g(\zeta_{n-1}^j, Y_{n-1}(\omega))}$,

Report $\hat{Z}_n^{\omega,N} = \hat{Z}_{n-1}^{\omega,N} \cdot \frac{\sum_{i=1}^N \tilde{g}^{\theta^{n-1}\omega}(\zeta_{n-1}^i)}{\sum_{i=1}^N \psi^{\theta^n \omega}(\zeta_n^i)}$.

In algorithmic terms, the difference between the sampling steps of Algorithm 2 and Algorithm 1 is fairly subtle: loosely speaking, at each time step, $N - 1$ of the particles in Algorithm 2 are propagated by the same mechanism as in Algorithm 1. However, with an appropriate choice of ψ^ω , the fluctuation properties of $\hat{Z}_n^{\omega,N}$ under (12)-(13) can be dramatically different to those of $Z_n^{\omega,N}$ under (8)-(7).

Our third main result (Theorem 2) concerns asymptotic fluctuation properties of twisted particle approximations in the regime where n and ω are fixed and $N \rightarrow \infty$. Under mild regularity conditions, we prove central limit theorems for generic particle systems under transitions like (12)-(13). For bounded test functions φ , we find that the $N \rightarrow \infty$ asymptotic variance associated with the empirical averages $N^{-1/2} \sum_{i=1}^N \varphi(\zeta_n^i)$ is the same when sampled under Algorithms 1 and 2, but the asymptotic variance associated with $\sqrt{N} \hat{Z}_n^{\omega,N}$ sampled Algorithm 1 is in general different to the asymptotic variance associated with $\sqrt{N} \hat{Z}_n^{\omega,N}$ sampled under Algorithm 2.

The finite- N , finite n behaviour of the relative variance of the standard estimate $Z_n^{\omega,N}$ from Algorithm 1 is well understood. Under certain regularity assumptions, it can be deduced from existing theory for standard particle algorithms [C  rou et al., 2011, Theorem 5.1] that in our setting $\Upsilon_N(\mathbf{M})$ must satisfy

$$\Upsilon_N(\mathbf{M}) \leq \log \left[1 + \frac{C}{N-1} \right], \quad (14)$$

for some finite constant C which depends on g and f . Our fourth main result (Proposition 5) generalizes (14) to the case of twisted particle filters. With $\Upsilon_N(\widetilde{\mathbf{M}})$ as in (10) and $\widetilde{\mathbf{M}}^\omega$ as in (12), application of Proposition 5 shows that

$$\Upsilon_N(\widetilde{\mathbf{M}}) \leq \log \left[1 + \frac{\mathcal{D}_{\mathbb{P}}(\psi, h)}{N-1} \right],$$

where

$$\mathcal{D}_{\mathbb{P}}(\psi, h) := \mathbb{P} - \text{ess sup}_\omega \left\{ \left[2 \sup_{z, z' \in \mathbf{X}} \frac{\psi^\omega(z)}{\psi^\omega(z')} - 1 \right] \sup_{z \in \mathbf{X}} \left(\frac{\psi^\omega(z)}{h^\omega(z)} \right) \sup_{(z, z') \in \mathbf{X}^2} \left| \frac{h^\omega(z)}{\psi^\omega(z)} - \frac{h^\omega(z')}{\psi^\omega(z')} \right| \right\}.$$

Thus whenever $\Upsilon_N(\mathbf{M}) > 0$, by choosing ψ “close” to h so that the oscillations of h^ω/ψ^ω are controlled, we can, in principle, achieve $\Upsilon_N(\widetilde{\mathbf{M}}) < \Upsilon_N(\mathbf{M})$.

Some remarks on connections to the literature are in order. Part of the generalized eigen-value theory through which h^ω arises (Proposition 2) is a variant on the theme of Kifer's Perron-Frobenius theorem for positive operators in a random environment [Kifer, 1996, Theorem 3.1], our conditions and technique of proof are different. Re-weighted particle transitions like (12) are reminiscent of certain eigen-function transformations of general-type branching processes studied by Athreya [2000], and more broadly can be viewed as a randomized version of Doob's h -transform. To the knowledge of the authors, such transformations applied to the type particle systems we consider have not previously been studied, or exploited for algorithmic purposes. Lastly, part of the proof of our Theorem 1 generalizes the proof of the necessity part in [Bucklew et al., 1990, Theorem 3], to the case of families of kernels driven an ergodic measure preserving transform.

The rest of the paper is structured as follows. Section 2 introduces our general setting, addresses the generalized eigen-value properties of families of non-negative kernels and sampling laws of the particle systems we consider. Section 3 narrows attention to a particular class of particle transitions, which we refer to as "twisted" and consider some of their properties in the regime where N is fixed, and $n \rightarrow \infty$, and vice-versa. Section 4 discusses the application of our main results to sequential importance sampling, bootstrap and auxiliary particle filters. Most of the proofs are given in section 5.

2 Non-negative kernels, sampling particles and variance growth

2.1 Notation and conventions

Henceforth we fix two measurable spaces $(X, \mathcal{X})$ and $(Y, \mathcal{Y})$. We are going to investigate a class of particle algorithms whose proposal and resampling mechanisms at each algorithmic time step may depend on an arbitrarily large number of observations, in quite a general fashion, and we will study properties of these algorithms under some assumptions of stationarity and ergodicity of the observation process. To this end let $\Omega = Y^{\mathbb{Z}}$ be the set of doubly infinite sequences valued in Y , let $\mathcal{F} = \mathcal{Y}^{\otimes \mathbb{Z}}$ and let $Y = \{Y_n\}_{n \in \mathbb{Z}}$ be the coordinate process on Ω , i.e. $Y_n(\omega) = \omega(n)$ for $\omega = \{\omega(n)\}_{n \in \mathbb{Z}} \in \Omega$. Let $\theta : \Omega \rightarrow \Omega$ be the shift operator $(\theta\omega)(n) := \omega(n+1)$, $n \in \mathbb{Z}, \omega \in \Omega$. For $n \in \mathbb{Z}$, the n -fold iterate of θ is written θ^n , and we take $\theta^0 := Id$. Let $\mathbb{P}$ be a probability measure on $(\Omega, \mathcal{F})$. Expectation w.r.t. $\mathbb{P}$ will be denoted by $\mathbb{E}$.

Let $\mathcal{M}(X)$, $\mathcal{P}(X)$ and $\mathcal{L}(X)$ be respectively the collections of measures, probability measures and bounded, real-valued, $\mathcal{X}$ -measurable functions on X . We write

$$\|\varphi\| := \sup_x |\varphi(x)|$$

and

$$\mu(\varphi) := \int_X \varphi(x) \mu(dx), \quad \text{for any } \varphi \in \mathcal{L}(X), \mu \in \mathcal{M}(X). \quad (15)$$

Whenever discussing functions whose co-domain is $\mathbb{R}$, we will assume that $\mathbb{R}$ is equipped with the Borel σ -algebra and all measurability statements regarding such functions will implicitly pertain to this σ -algebra on their co-domain.

We will be dealing throughout with various real-valued functions on $\Omega \times X$ (and more generally $\Omega \times X^N$ etc.). For any such function φ , we write the ω -section of φ as $\varphi^\omega : X \rightarrow \mathbb{R}$, $\varphi^\omega(x) := \varphi(\omega, x)$. For a function $\xi : \Omega \rightarrow \mathbb{R}$ it will sometimes be convenient to write ξ_ω instead of the more standard $\xi(\omega)$.

We will need to express various integration operations involving functions on $\Omega \times X^N$ and their ω -sections, so for completeness we quote the following facts of measure theory, see for example [Doob, 1994, Chapter VI], which will be used in the sequel in various places without further comment: when $\varphi : \Omega \times X^N \rightarrow \mathbb{R}$ is measurable w.r.t. to $\mathcal{F} \otimes \mathcal{X}^{\otimes N}$, then for every $\omega \in \Omega$, the ω -section φ^ω , is measurable w.r.t. $\mathcal{X}^{\otimes N}$; and furthermore for any σ -finite measure μ on $(X^N, \mathcal{X}^{\otimes N})$, if φ is integrable w.r.t. to $\mathbb{P} \otimes \mu$, then the function acting $\Omega \rightarrow \mathbb{R}$ which maps $\omega \mapsto \mu(\varphi^\omega)$ is measurable w.r.t. $\mathcal{F}$ and is $\mathbb{P}$ -integrable.

Let $N \geq 1$ be fixed and let $\varphi, \tilde{\varphi}$ be two functions, each acting $\Omega \times X^N \rightarrow \mathbb{R}$ and each measurable w.r.t. $\mathcal{F} \otimes \mathcal{X}^{\otimes N}$. We will need to talk about the sets on which two such functions take the same values. For any $\omega \in \Omega$, let $A_\omega := \{x \in X^N : \varphi^\omega(x) = \tilde{\varphi}^\omega(x)\}$ and let μ be a σ -finite measure on $(X^N, \mathcal{X}^{\otimes N})$. In order to avoid having to make the sets $\{A_\omega; \omega \in \Omega\}$ explicit in various statements, we will write by convention:

$$\text{for } \mathbb{P} - a.a. \omega, \quad \varphi^\omega(x) = \tilde{\varphi}^\omega(x), \quad \text{for } \mu - a.a. x$$

to mean

$$\mathbb{P}(\{\omega : \mu(A_\omega^c) = 0\}) = 1.$$

2.2 Generalized eigen-value theory for non-negative kernels

Fix arbitrarily $\mu_0 \in \mathcal{P}(\mathsf{X})$ and let $M : \Omega \times \mathsf{X} \times \mathcal{X} \rightarrow [0, 1]$ be such that $M(\omega, x, \cdot) \in \mathcal{P}(\mathsf{X})$ for each $(\omega, x) \in \Omega \times \mathsf{X}$, and $M(\cdot, \cdot, A)$ is $\mathcal{F} \otimes \mathcal{X}$ -measurable for each $A \in \mathcal{X}$. Then for any ω , $M(\omega, \cdot, \cdot)$ is a Markov kernel on $(\mathsf{X}, \mathcal{X})$ and when it is important to emphasize this perspective, we shall often write $M^\omega(x, A)$ instead of $M(\omega, x, A)$. We shall adopt similar notation for other kernels.

For any fixed $\omega \in \Omega$, let $\mathbb{E}^\omega$ denote expectation with respect to the law of the time-inhomogeneous Markov chain $\{X_n; n \geq 0\}$, with each X_n valued in X , initialized from $X_0 \sim \mu_0$ and $X_n | \{X_{n-1} = x_{n-1}\} \sim M^{\theta^{n-1}\omega}(x_{n-1}, \cdot)$, for $n \geq 1$. Let $G : \Omega \times \mathsf{X} \rightarrow \mathbb{R}_+$ be a $\mathcal{F} \otimes \mathcal{X}$ -measurable, strictly positive and bounded function.

Remark 1. This setup is purposefully generic and accommodates, as one particular instance, the case

$$G^\omega(x) = g(x, Y_0(\omega)), \quad M^\omega(x, dx') = f(x, dx'), \quad \forall \omega \in \Omega. \quad (16)$$

where g and f are as in section 1, and then

$$\mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p) \right] \equiv Z_n^\omega.$$

Furthermore, with the particular G, M as in (16), the generic particle system which we will go on to introduce in section 2.3 is the bootstrap particle filter discussed in section 1. However, we stress that (16) is just one instance of our generic setup, others will be discussed in section 4.

We next introduce two hypotheses.

(H1) The shift operator θ preserves $\mathbb{P}$ and is ergodic.

(H2) There exist constants $\beta \in [1, \infty)$, $(\epsilon_-, \epsilon_+) \in (0, \infty)^2$, and $\nu \in \mathcal{P}(\mathsf{X})$ such that

$$\frac{G(\omega, x)}{G(\omega', x')} \leq \beta, \quad \forall (\omega, \omega', x, x') \in \Omega^2 \times \mathsf{X}^2, \quad (17)$$

$$\epsilon_- \nu(\cdot) \leq M(\omega, x, \cdot) \leq \epsilon_+ \nu(\cdot), \quad \forall (\omega, x) \in \Omega \times \mathsf{X}. \quad (18)$$

Since in our setup we have taken $\Omega := \mathsf{Y}^\mathbb{Z}$, hypothesis **(H1)** amounts to saying that the observation process is stationary and ergodic. However, the setup $\Omega := \mathsf{Y}^\mathbb{Z}$ has only been chosen in order to connect out general framework with the motivating context of HMMs and thereby provide intuition to the reader; in order for **(H1)** to be meaningful and the arguments in the proofs of our main results to hold, it is enough that $(\Omega, \mathcal{F}, \mathbb{P})$ is just some abstract probability space, and that $\theta : \Omega \rightarrow \Omega$ is an invertible transform.

Hypothesis **(H2)** is a strong mixing condition, and will rarely hold when X and Y are non-compact. We adopt this hypothesis partly for convenience of presentation; in some places in our proofs for the results of section 2.2 where we adopt **(H2)**, our arguments will not actually rely on both (17) and (18) holding simultaneously, but we package these conditions together in order to avoid a layer of technical presentation which would further lengthen and complicate our proofs, our priority in section 2.2 is to move swiftly through these preparatory details and on to the more novel matter which appears later. Some further remarks on this issue are suppressed until the end of section 2.2.

It is well known that **(H1)** and **(H2)** are together more than enough to establish the following result - see [Leroux, 1992] for related ideas in the context of HMMs - and we include a proof only for purposes of exposition: the phenomena it describes and some of the objects involved are the cornerstone for what follows in subsequent sections.

Proposition 1. Assume **(H1)** and **(H2)**. Then there exists a constant $\Lambda \in (-\infty, \infty)$ independent of the initial distribution $\mu_0 \in \mathcal{P}(\mathbf{X})$ such that

$$\frac{1}{n} \log \mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p) \right] \rightarrow \Lambda, \quad \text{as } n \rightarrow \infty, \quad \mathbb{P} - a.s. \quad (19)$$

The proof of Proposition 1, given below, can be expressed conveniently using a little further operator-type notation. We now introduce the non-negative kernel

$$Q : \Omega \times \mathbf{X} \times \mathcal{X} \rightarrow \mathbb{R}_+, \quad Q(\omega, x, dx') := G(\omega, x)M(\omega, x, dx').$$

For any fixed $\omega \in \Omega$ define the operators

$$Q^\omega(\varphi)(x) := \int_{\mathbf{X}} Q^\omega(x, dx') \varphi(x'), \quad \varphi \in \mathcal{L}(\mathbf{X}), \quad (20)$$

$$\mu Q^\omega(\cdot) := \int_{\mathbf{X}} \mu(dx) Q^\omega(x, \cdot), \quad \mu \in \mathcal{M}(\mathbf{X}). \quad (21)$$

and let $\{Q_n^\omega; n \in \mathbb{N}\}$ be defined inductively by

$$Q_0^\omega := Id, \quad Q_n^\omega = Q_{n-1}^\omega Q^{\theta^{n-1} \omega}, \quad n \geq 1. \quad (22)$$

This operator notation allows us to express

$$\mu_0 Q_n^\omega(1) = \mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p) \right], \quad n \geq 1. \quad (23)$$

Proof. (of Proposition 1). Under **(H2)** we have $\underline{g} := \inf_{\omega, x} G^\omega(x) > 0$. Now consider the sequence of random variables $\{\kappa_n^\omega; n \geq 1\}$ defined by

$$\kappa_n^\omega := \nu Q_{n-1}^\omega(1) \underline{g} \epsilon_-.$$

From the definition of (22) it is straightforward to establish by induction the following semigroup property: for any $\omega \in \Omega$, and $p, n \geq 0$,

$$Q_{p+n}^\omega = Q_p^\omega Q_n^{\theta^p \omega},$$

and this combined with **(H2)** gives

$$\begin{aligned} \kappa_{p+n-1}^\omega &= \nu Q_p^\omega Q_{n-1}^{\theta^p \omega}(1) \underline{g} \epsilon_- \\ &= \nu Q_{p-1}^\omega Q^{\theta^{p-1} \omega} Q_{n-1}^{\theta^p \omega}(1) \underline{g} \epsilon_- \\ &\geq \kappa_p^\omega \kappa_n^{\theta^p \omega}, \end{aligned}$$

and so

$$-\log \kappa_{p+n}^\omega \leq -\log \kappa_p^\omega - \log \kappa_n^{\theta^p \omega}.$$

Furthermore under **(H2)**, $\sup_{\omega, x} G^\omega(x) < \infty$, so there exists a finite constant c such that

$$\int_{\Omega} -\log \kappa_n^\omega \mathbb{P}(d\omega) \geq -cn$$

for any $n \geq 1$, and these considerations, combined **(H1)**, allow application of Kingman's sub-additive ergodic theorem to establish that there exists a constant $\Lambda \in (-\infty, \infty)$ such that

$$\frac{1}{n} \log \kappa_n^\omega \longrightarrow \Lambda, \quad \text{as } n \rightarrow \infty, \quad \mathbb{P} - a.s.$$

Under **(H2)**, for any $\omega \in \Omega$, $n \geq 1$, $x \in \mathbf{X}$ and $\mu \in \mathcal{P}(\mathbf{X})$,

$$\frac{\nu Q_n^\omega(1)}{\mu Q_n^\omega(1)} = \frac{\nu Q^\omega Q_{n-1}^{\theta \omega}(1)}{\mu Q^\omega Q_{n-1}^{\theta \omega}(1)} \leq \beta \frac{\epsilon_+}{\epsilon_-},$$

and combining this with a lower bound of a similar form we find

$$\sup_{\omega \in \Omega} \left| \frac{1}{n} \log \kappa_n^\omega - \frac{1}{n} \log \mu Q_n^\omega(1) \right| \leq \frac{1}{n} \log \left(\beta \frac{\epsilon_+}{\epsilon_-} \right) + \frac{1}{n} |\log(\epsilon_- \underline{g})|$$

so

$$\frac{1}{n} \log \mu Q_n^\omega(1) \longrightarrow \Lambda, \quad \text{as } n \rightarrow \infty \quad \mathbb{P} - a.s.$$

The proof is complete upon noting (23). $\square$

It turns out that Proposition 1 is one element of a generalized eigen-value theory for the non-negative kernel Q . Under **(H1)** and **(H2)**, Proposition 2 below, directly inspired by [Kifer, 1996, Theorem 3.1], identifies a function $h : \Omega \times \mathbf{X} \rightarrow \mathbb{R}_+$ and a random variable $\lambda : \Omega \times \mathbf{X} \rightarrow \mathbb{R}_+$ satisfying

$$Q^\omega(h^{\theta^\omega}) = \lambda_\omega h^\omega.$$

As we shall go on to see, the expected value of $\log \lambda$ under $\mathbb{P}$ is equal to the constant Λ appearing in Proposition 1 and h plays a special role in the construction of efficient particle algorithms.

Now let $\Phi^\omega : \mathcal{P}(\mathbf{X}) \rightarrow \mathcal{P}(\mathbf{X})$ be defined by

$$\Phi^\omega(\mu) = \frac{\mu Q^\omega}{\mu Q^\omega(1)}, \quad \mu \in \mathcal{P}(\mathbf{X}),$$

and let $\{\Phi_n^\omega; n \in \mathbb{N}\}$ be the family of operators defined inductively by

$$\Phi_0^\omega := Id, \quad \Phi_n^\omega := \Phi^{\theta^{n-1}\omega} \circ \Phi_{n-1}^\omega,$$

so that each Φ_n^ω acts $\mathcal{P}(\mathbf{X}) \rightarrow \mathcal{P}(\mathbf{X})$. Under these definitions, for any $n \in \mathbb{N}$,

$$\Phi_n^\omega(\mu) = \frac{\mu Q_n^\omega}{\mu Q_n^\omega(1)}, \tag{24}$$

which can be verified by induction, since from the above definitions $\Phi_0^\omega = Id$, $Q_0^\omega := Id$ and when (24) holds,

$$\Phi_{n+1}^\omega(\mu) = \left(\Phi^{\theta^n \omega} \circ \Phi_n^\omega \right) (\mu) = \frac{\Phi_n^\omega(\mu) Q^{\theta^n \omega}}{\Phi_n^\omega(\mu) Q^{\theta^n \omega}(1)} = \frac{\mu Q_n^\omega Q^{\theta^n \omega}}{\mu Q_n^\omega Q^{\theta^n \omega}(1)} = \frac{\mu Q_{n+1}^\omega}{\mu Q_{n+1}^\omega(1)}.$$

Remark 2. In the setting $M^\omega(x, dx') := f(x, dx')$, $G^\omega(x) := g(x, Y(\omega))$, then if μ_0 and π_n^ω are respectively the initial distribution and prediction-filter as in (2), we have

$$\pi_{n+1}^\omega = \Phi^{\theta^n \omega}(\pi_n^\omega), \quad n \geq 0.$$

Remark 3. Under the uniform bounds of **(H2)**, it is known that Φ_n^ω is exponentially stable with respect to initial conditions [e.g., Del Moral, 2004, Chapter 4] in the sense there exists constants $C < \infty$ and $\rho < 1$ such that for any $\varphi \in \mathcal{L}(\mathbf{X})$ and any $n \geq 1$,

$$\sup_{\omega \in \Omega} \sup_{\mu, \mu' \in \mathcal{P}(\mathbf{X})} |[\Phi_n^\omega(\mu) - \Phi_n^\omega(\mu')](\varphi)| \leq \|\varphi\| C \rho^n. \tag{25}$$

(25) is used extensively in the proof of the following proposition, given in section 5.

Proposition 2. Assume **(H2)**.

1) Fix $\mu \in \mathcal{P}(\mathbf{X})$. Then the limits

$$\eta^\omega(A) := \lim_{n \rightarrow \infty} \Phi_n^{\theta^{-n}\omega}(\mu)(A), \quad \omega \in \Omega, \quad A \in \mathcal{X}, \tag{26}$$

$$h(\omega, x) := \lim_{n \rightarrow \infty} \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\mu) Q_n^\omega(1)}, \quad \omega \in \Omega, \quad x \in \mathbf{X}, \tag{27}$$

exist and define a family of probability measures $\eta := \{\eta^\omega \in \mathcal{P}(\mathbf{X}); \omega \in \Omega\}$, and an $\mathcal{F} \otimes \mathcal{X}$ -measurable function $h : \Omega \times \mathbf{X} \rightarrow \mathbb{R}$.

2) In fact, η and h are independent of the particular μ chosen in part 1) and there exist constants $C < \infty$ and $\rho < 1$ such that for any $\varphi \in \mathcal{L}(\mathbf{X})$,

$$\sup_{\omega \in \Omega} \sup_{\mu \in \mathcal{P}(\mathbf{X})} \left| \left[\Phi_n^{\theta^{-n}\omega}(\mu) - \eta^\omega \right] (\varphi) \right| \leq \|\varphi\| C \rho^n, \quad n \geq 1, \quad (28)$$

and

$$\sup_{\omega \in \Omega} \sup_{x \in \mathbf{X}} \sup_{\mu \in \mathcal{P}(\mathbf{X})} \left| \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\mu) Q_n^\omega(1)} - h(\omega, x) \right| \leq C \rho^n, \quad n \geq 1. \quad (29)$$

3) $\lambda : \omega \in \Omega \mapsto \eta^\omega(G^\omega) \in \mathbb{R}_+$ is measurable w.r.t. $\mathcal{F}$ and we have the bounds

$$\sup_{(\omega, \omega') \in \Omega^2} \frac{\lambda_\omega}{\lambda_{\omega'}} < \infty, \quad \sup_{(\omega, \omega', x, x') \in \Omega^2 \times \mathbf{X}^2} \frac{h(\omega, x)}{h(\omega', x')} < \infty. \quad (30)$$

4) Amongst all triples which consist of: (i) an Ω -indexed family of probability measures on $(\mathbf{X}, \mathcal{X})$, (ii) an $\mathbb{R}_+$ -valued, not identically zero, measurable function on $\Omega \times \mathbf{X}$, and (iii) a measurable function on Ω ; the triple (η, h, λ) , with η, h as in part 1) and λ as in part 3), uniquely satisfies the system of equations

$$\eta^\omega Q^\omega = \lambda_\omega \eta^{\theta\omega}, \quad Q^\omega(h^{\theta\omega}) = \lambda_\omega h^\omega, \quad \eta^\omega(h^\omega) = 1, \quad \text{for all } \omega \in \Omega. \quad (31)$$

The connection with Proposition 1 is:

Proposition 3. Assume **(H1)**, **(H2)** and let Λ be as in Proposition 1 and λ be as in Proposition 2. Then

$$\Lambda = \mathbb{E}[\log \lambda] = \int_{\Omega} \frac{Q^\omega(h^{\theta\omega})(x)}{h^\omega(x)} \mathbb{P}(d\omega), \quad \text{for any } x \in \mathbf{X}. \quad (32)$$

In the setting of HMMs as per Remark 1, equalities like the first one in (32) appear routinely in the study of likelihood-based estimators [Leroux, 1992, Douc and Moulines, 2011]. However, it is the second equality in (32), and generalizations thereof, which shall be crucial for our purposes in the sequel.

Remark 4. If one weakens the “1-step” condition (18) to an m -step version for some $m \geq 1$, then Propositions 1-3 can easily be generalized, since one can simply work with the kernel Q_m^ω instead of Q^ω . The utility of imposing the uniform in ω and x bounds in **(H2)** is that the statements of Proposition 2 hold for all $\omega \in \Omega$, and without having to invoke **(H1)**. If one allows ω -dependent constants and measures in (17) and (18), but instead imposes certain explicit compactness continuity assumptions and **(H1)**, then [Kifer, 1996, Theorem 3.1] provides a partial alternative to our Proposition 2.

We shall proceed by introducing the laws of the particle systems of interest.

2.3 Law of the standard particle system

Unless stated otherwise, in this section we fix arbitrarily $N \geq 1$ and write $\mathcal{P}(\mathbf{X}^N)$ for the collection of probability measures on $(\mathbf{X}^N, \mathcal{X}^{\otimes N})$.

Let $\mathbf{M} : \Omega \times \mathbf{X}^N \times \mathcal{X}^{\otimes N} \rightarrow [0, 1]$ be given, in integral form, by

$$\mathbf{M}(\omega, x, dz) = \prod_{i=1}^N \left[\frac{\sum_{j=1}^N G(\omega, x^j) M(\omega, x^j, dz^i)}{\sum_{j=1}^N G(\omega, x^j)} \right], \quad (33)$$

where $x = (x^1, \dots, x^N)$, $z = (z^1, \dots, z^N) \in \mathbf{X}^N$. Each member of the family $\{\mathbf{M}^\omega; \omega \in \Omega\}$ is a Markov transition kernel for the entire N -particle system according to a “multinomial” resampling scheme with fitness function $G(\omega, \cdot)$, followed by conditionally independent mutation according to M^ω .

Now for any given $\omega \in \Omega$, we shall denote by $\mathbb{E}_N^\omega$ expectation with respect to the law of the Markov chain $\{\zeta_n; n \geq 0\}$, with each $\zeta_n = \{\zeta_n^1, \dots, \zeta_n^N\}$ valued in $\mathbf{X}^N$ and

$$\zeta_0 \sim \mu_0^{\otimes N}, \quad \zeta_n | \zeta_{n-1} \sim \mathbf{M}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot). \quad (34)$$

Define

$$\mathbf{G} : (\omega, x) \in \Omega \times \mathbf{X}^N \mapsto \frac{1}{N} \sum_{i=1}^N G(\omega, x^i) \in \mathbb{R}_+,$$

where $x = (x^1, \dots, x^N)$.

Remark 5. For any $\varphi \in \mathcal{L}(\mathbf{X})$, if we define the function

$$\varphi : x = (x^1, \dots, x^N) \in \mathbf{X}^N \mapsto \frac{1}{N} \sum_{i=1}^N \varphi(x^i) \in \mathbb{R},$$

then the well known lack-of-bias property of the particle approximation reads as

$$\mathbb{E}_N^\omega \left[\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p \omega}(\zeta_p) \varphi(\zeta_n) \right] = \mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p) \varphi(X_n) \right]. \quad (35)$$

Remark 6. When $M^\omega(x, \cdot) = f(x, \cdot)$ and $G^\omega(x) = g(x, Y_0(\omega))$, the sampling recipe for simulating the process $\{\zeta_n; n \geq 0\}$ according to (34) is the bootstrap particle filter: algorithm 1. Furthermore, the particle approximation of Z_n^ω is then given by

$$\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p \omega}(\zeta_p).$$

Part of our investigation will be to develop some limit theory for

$$\frac{\mathbb{E}_N^\omega \left[\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p \omega}(\zeta_p)^2 \right]}{\mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(\zeta_p) \right]^2} \quad (36)$$

when N is fixed and $n \rightarrow \infty$. Our notation $\mathbf{G}, \mathbf{M}$ and the display (35) are intended to hint that the type of arguments employed in the proof of Proposition 1 and the phenomena described in Proposition 2 may be relevant to the study of (36). Indeed this is the direction in which we are heading. However we will actually study an object more general than (36), arising from a more general form of particle approximation, which we now introduce. The generality arises as we consider the particle system distributed according to some Markov law, possibly different to (34).

2.4 Alternative sampling of the particle system

Let us introduce $\widetilde{\mathbf{M}} : \Omega \times \mathbf{X}^N \times \mathcal{X}^{\otimes N} \rightarrow [0, 1]$, possibly different from $\mathbf{M}$. For fixed ω , now denote by $\widetilde{\mathbb{E}}_N^\omega$ expectation with respect to law of the Markov chain

$$\zeta_0 \sim \mu_0^{\otimes N}, \quad \zeta_n | \zeta_{n-1} \sim \widetilde{\mathbf{M}}^{\theta^{n-1} \omega}(\zeta_{n-1}, \cdot). \quad (37)$$

We are going to specify a class of candidates for $\widetilde{\mathbf{M}}$, and we first notice that the regularity condition **(H2)** transfers to $\mathbf{G}, \mathbf{M}$ in the following sense (the proof is elementary):

Lemma 1. *Assume **(H2)**. Then for any $N \geq 1$,*

$$\frac{\mathbf{G}(\omega, x)}{\mathbf{G}(\omega', x')} \leq \beta, \quad \forall (\omega, \omega', x, x') \in \Omega \times \mathbf{X}^{2N},$$

$$\frac{1}{\beta} \epsilon_-^N \nu^{\otimes N}(\cdot) \leq \mathbf{M}(\omega, x, \cdot) \leq \beta \epsilon_+^N \nu^{\otimes N}(\cdot), \quad \forall (\omega, x) \in \Omega \times \mathbf{X}^N.$$

We shall consider the following family of kernels.

Definition. ($\mathbb{M}$) Any $\widetilde{\mathbf{M}} : \Omega \times \mathbf{X}^N \times \mathcal{X}^{\otimes N} \rightarrow [0, 1]$ is a member of $\mathbb{M}$ if and only if there exist constants $(\widetilde{\epsilon}_-, \widetilde{\epsilon}_+) \in (0, \infty)^2$ and $\widetilde{\nu} \in \mathcal{P}(\mathbf{X}^N)$ such that

$$\widetilde{\nu}(\cdot) \widetilde{\epsilon}_- \leq \widetilde{\mathbf{M}}(\omega, x, \cdot) \leq \widetilde{\epsilon}_+ \widetilde{\nu}(\cdot), \quad \forall (\omega, x) \in \Omega \times \mathbf{X}^N, \quad \nu^{\otimes N} \ll \widetilde{\nu}, \quad \text{and} \quad \int_{\mathbf{X}^N} \left(\frac{d\nu^{\otimes N}}{d\widetilde{\nu}}(x) \right)^2 \widetilde{\nu}(dx) < \infty, \quad (38)$$

where ν is as in **(H2)**.

When $\widetilde{\mathbf{M}}$ is a member of $\mathbb{M}$ we write

$$\phi^\omega(x, x') := \frac{d\mathbf{M}^\omega(x, \cdot)}{d\widetilde{\mathbf{M}}^\omega(x, \cdot)}(x'), \quad (\omega, x, x') \in \Omega \times \mathbf{X}^{2N}, \quad (39)$$

and in the context of sampling the particle system $(\zeta_n; n \geq 0)$ under the law (37), we will take

$$\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p \omega}(\zeta_p) \phi^{\theta^p \omega}(\zeta_p, \zeta_{p+1}) \quad (40)$$

as an approximation of $\mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p) \right]$. Clearly (40) is unbiased under (37), in the sense that

$$\widetilde{\mathbb{E}}_N^\omega \left[\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p \omega}(\zeta_p) \phi^{\theta^p \omega}(\zeta_p, \zeta_{p+1}) \right] = \mathbb{E}_N^\omega \left[\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p \omega}(\zeta_p) \right] = \mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p) \right],$$

where the first equality is immediate under (37) and (40), and the second equality is the lack-of-bias property as per Remark 5. The following result describes the $n \rightarrow \infty$ behaviour of

$$\widetilde{\mathcal{V}}_{n,N}^\omega := \frac{\widetilde{\mathbb{E}}_N^\omega \left[\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p \omega}(\zeta_p)^2 \phi^{\theta^p \omega}(\zeta_p, \zeta_{p+1})^2 \right]}{\mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p) \right]^2}. \quad (41)$$

Its proof, provided in section 5, starts by considering the family of kernels $\{\widetilde{\mathbf{R}}^\omega; \omega \in \Omega\}$, with

$$\widetilde{\mathbf{R}}^\omega(x, dx') := \mathbf{G}^\omega(x)^2 \phi^\omega(x, x')^2 \widetilde{\mathbf{M}}^\omega(x, dx'),$$

in terms of which the numerator of (41) may be written, and which exhibit exactly similar properties to those of Q^ω appearing in the proof of Proposition (1).

Proposition 4. Assume **(H1)**, **(H2)** and fix $N \geq 1$ arbitrarily. For every $\widetilde{\mathbf{M}} \in \mathbb{M}$ there exists a constant $\Upsilon_N(\widetilde{\mathbf{M}}) \in [0, \infty)$, independent of the initial distribution μ_0 such that

$$\frac{1}{n} \log \widetilde{\mathcal{V}}_{n,N}^\omega \longrightarrow \Upsilon_N(\widetilde{\mathbf{M}}), \quad \text{as } n \rightarrow \infty, \quad \mathbb{P} - a.s.$$

We now proceed to address the question of how $\Upsilon_N(\widetilde{\mathbf{M}})$ depends on $\widetilde{\mathbf{M}}$. To this end, let us introduce two further pieces of notation:

$$\mathbf{Q}(\omega, x, dx') := \mathbf{G}(\omega, x) \mathbf{M}(\omega, x, dx'),$$

and when **(H2)** holds, so that h as in Proposition 2 is well-defined, consider the function

$$\mathbf{h} : (\omega, x) \in \Omega \times \mathbf{X}^N \longmapsto \frac{1}{N} \sum_{i=1}^N h(\omega, x^i) \in \mathbb{R}_+. \quad (42)$$

Our interest in (42) stems from the following pivotal lemma, which shows how the eigen-function h and eigen-value λ of Q appearing in Proposition 2 define an eigen-function and eigen-value for $\mathbf{Q}$. Recall that $N \geq 1$ is still arbitrarily fixed.

Lemma 2. For any $\omega \in \Omega$,

$$\mathbf{Q}^\omega(\mathbf{h}^{\theta\omega}) = \lambda_\omega \mathbf{h}^\omega,$$

where λ_ω is as in Proposition 2.

Proof.

$$\begin{aligned} \mathbf{Q}^\omega(\mathbf{h}^{\theta\omega})(x) &= \frac{1}{N} \sum_{k=1}^N \int_{\mathbf{X}^N} \mathbf{Q}^\omega(x, dz) h^{\theta\omega}(z^k) \\ &= \frac{1}{N} \sum_{k=1}^N \mathbf{G}^\omega(x) \int_{\mathbf{X}} \frac{\sum_{i=1}^N Q^\omega(x^i, dz^k)}{\sum_{i=1}^N G^\omega(x^i)} h^{\theta\omega}(z^k) \\ &= \frac{1}{N} \sum_{k=1}^N \frac{1}{N} \sum_{i=1}^N \int_{\mathbf{X}} Q^\omega(x^i, dz^k) h^{\theta\omega}(z^k) \\ &= \lambda_\omega \frac{1}{N} \sum_{i=1}^N h^\omega(x^i) = \lambda_\omega \mathbf{h}^\omega(x). \end{aligned}$$

□

Now consider taking

$$\widetilde{\mathbf{M}}^\omega(x, dx') = \frac{\mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}, \quad (43)$$

which is clearly a member of $\mathbb{M}$, due to the definition of $\mathbf{h}$ and part 4) of Proposition 2. In this case we have

$$\begin{aligned} \prod_{p=0}^{n-1} \mathbf{G}^{\theta^p\omega}(\zeta_p) \phi^{\theta^p\omega}(\zeta_p, \zeta_{p+1}) &= \prod_{p=0}^{n-1} \mathbf{G}^{\theta^p\omega}(\zeta_p) \frac{\int_{\mathbf{X}^N} \mathbf{M}^{\theta^p\omega}(\zeta_p, dz_{p+1}) \mathbf{h}^{\theta^{p+1}\omega}(z_{p+1})}{\mathbf{h}^{\theta^{p+1}\omega}(\zeta_{p+1})} \\ &= \prod_{p=0}^{n-1} \frac{\mathbf{Q}^{\theta^p\omega}(\mathbf{h}^{\theta^{p+1}\omega})(\zeta_p)}{\mathbf{h}^{\theta^{p+1}\omega}(\zeta_{p+1})} \\ &= \frac{\mathbf{h}^\omega(\zeta_0)}{\mathbf{h}^{\theta^n\omega}(\zeta_n)} \prod_{p=0}^{n-1} \frac{\mathbf{Q}^{\theta^p\omega}(\mathbf{h}^{\theta^{p+1}\omega})(\zeta_p)}{\mathbf{h}^{\theta^{p+1}\omega}(\zeta_{p+1})} \\ &= \frac{\mathbf{h}^\omega(\zeta_0)}{\mathbf{h}^{\theta^n\omega}(\zeta_n)} \prod_{p=0}^{n-1} \lambda_{\theta^p\omega}, \end{aligned} \quad (44)$$

where the final equality is due to Lemma 2. Thus if we choose $\widetilde{\mathbf{M}}$ as per (43), then the quantity in (44) depends on the particle system trajectory $\zeta_0, \dots, \zeta_n$ only through the quantities $\mathbf{h}^\omega(\zeta_0)$ and $\mathbf{h}^{\theta^n\omega}(\zeta_n)$, and we then might hope that $\Upsilon_N(\widetilde{\mathbf{M}}) = 0$. This turns out to be true, and much more strikingly, up to its definition on certain sets of measure zero, $\widetilde{\mathbf{M}}$ as in (43) is the unique member of $\mathbb{M}$ which achieves $\Upsilon_N(\widetilde{\mathbf{M}}) = 0$, in the sense of the following theorem.

Theorem 1. Assume (H1), (H2), let $N \geq 1$ be fixed arbitrarily and let $\widetilde{\mathbf{M}}$ be any member of $\mathbb{M}$. Then 1)-3) are equivalent.

$$1) \quad \Upsilon_N(\widetilde{\mathbf{M}}) = 0.$$

$$2) \quad \text{For } \mathbb{P}\text{-almost all } \omega \in \Omega \text{ there exists } A_\omega \in \mathcal{X}^{\otimes N} \text{ such that } \nu^{\otimes N}(A_\omega^c) = 0 \text{ and for any } x \in A_\omega,$$

$$\widetilde{\mathbf{M}}^\omega(x, B) = \frac{\int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}, \quad \text{for all } B \in \mathcal{X}^{\otimes N}. \quad (45)$$

$$3) \quad \text{For } \mathbb{P}\text{-almost all } \omega \in \Omega, \sup_n \widetilde{\mathcal{V}}_{n,N}^\omega < \infty.$$

The proof of Proposition 4 can be found in section 5. Showing 2) $\Rightarrow$ 3) amounts to little more than equation (44). It is obvious that 3) $\Rightarrow$ 1). Showing 1) $\Rightarrow$ 2) is the challenging part.

The following Lemma serves to accompany Proposition 4 and Theorem 1, and provides necessary and sufficient conditions for $\Upsilon_N(\widetilde{\mathbf{M}}) = 0$ in the case of taking $\widetilde{\mathbf{M}} = \mathbf{M}$, i.e. the transitions of the standard particle system. The proof is given in section 5.

Lemma 3. *Assume (H1), (H2) and let $N \geq 1$ be fixed arbitrarily. Then 1)-3) are equivalent.*

$$1) \ \Upsilon_N(\mathbf{M}) = 0$$

$$2) \ \text{For } \mathbb{P} - \text{a.a. } \omega, \quad h^\omega(x) = 1, \text{ for } \nu - \text{a.a. } x.$$

$$3) \ \text{There exists a random variable } C : \Omega \rightarrow \mathbb{R}_+ \text{ such that}$$

$$\text{for } \mathbb{P} - \text{a.a. } \omega, \quad G^\omega(x) = C_\omega, \quad \text{for } \nu - \text{a.a. } x.$$

In situations of practical interest, point 3) of Lemma 3 is usually false, and then it must be the case that $\Upsilon_N(\mathbf{M}) > 0$. It then appears that a choice of $\widetilde{\mathbf{M}}$ which approximates the optimal transition, (45), may yield a provable performance advantage over $\mathbf{M}$, in the sense of achieving strict inequality $\Upsilon_N(\widetilde{\mathbf{M}}) < \Upsilon_N(\mathbf{M})$. This leads us to consider the class of particle algorithms treated in the next section.

3 Twisted particle algorithms

The form of the optimal transition kernel (45) suggests that we consider families of kernels arising from re-weighting of $\mathbf{M}^\omega(x, \cdot)$ by an additive, non-negative functional. In this section we will analyze particle algorithms arising from kernels of this general form. Let $\psi : \Omega \times \mathbf{X} \rightarrow \mathbb{R}_+$ be a strictly positive, bounded and measurable function and then define

$$\psi : (\omega, x) \in \Omega \times \mathbf{X}^N \quad \mapsto \quad \frac{1}{N} \sum_{i=1}^N \psi(\omega, x^i) \in \mathbb{R}_+.$$

For the purposes of this section, let us consider the following mild regularity assumption.

(H3) For each $\omega \in \Omega$, $\sup_x G^\omega(x) < \infty$ and $\sup_x \psi^\omega(x) < \infty$.

When **(H3)** holds the following Markov kernel is well-defined.

$$\widetilde{\mathbf{M}}^\omega(x, dx') = \frac{\mathbf{M}^\omega(x, dx') \psi^{\theta\omega}(x')}{\int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dz) \psi^{\theta\omega}(z)}. \quad (46)$$

We shall analyze particle approximations which arise from sampling under (46). Our motivation here is that we have in mind situations where ψ is chosen to be some approximation of h , assuming the latter exists. The kernel (46) accommodates the standard transition (33) (e.g. take $\psi^\omega = 1$) and the optimal transition identified in Theorem 1 (take $\psi^\omega = h^\omega$).

This section addresses two main objectives. Firstly, to validate the particle approximations delivered when sampling under (46), by analyzing some of their convergence and fluctuation properties in the regime where $N \rightarrow \infty$. Secondly, to provide an estimate of $\Upsilon_N(\widetilde{\mathbf{M}})$ which exhibits dependence on N and on the discrepancy between ψ^ω and h^ω .

Let us introduce a little more notation. Define, for each $\omega \in \Omega$,

$$\eta_0 := \mu_0, \quad \eta_n^\omega := \Phi^{\theta^{n-1}\omega}(\eta_{n-1}^\omega), \quad n \geq 1.$$

$$\widehat{\Phi}^\omega : \mathcal{P}(\mathbf{X}) \rightarrow \mathcal{P}(\mathbf{X}), \quad \widehat{\Phi}^\omega(\mu)(dx) := \frac{\mu Q^\omega(dx) \psi^\omega(x)}{\mu Q^\omega(\psi^\omega)}, \quad \mu \in \mathcal{P}(\mathbf{X}).$$

and also for each $N \geq 1$,

$$\Gamma_N^\omega : \mathcal{P}(\mathbf{X}) \rightarrow \mathcal{P}(\mathbf{X}), \quad \Gamma_N^\omega(\mu) := \frac{1}{N} \widehat{\Phi}^\omega(\mu) + \left(1 - \frac{1}{N}\right) \Phi^\omega(\mu), \quad \mu \in \mathcal{P}(\mathbf{X}).$$

$$\Gamma_{0,N}^\omega := Id, \quad \Gamma_{n,N}^\omega := \Gamma_N^{\theta^{n-1}\omega} \circ \dots \circ \Gamma_N^{\theta\omega} \circ \Gamma_N^\omega, \quad n \geq 1.$$

For purposes of developing limits and fluctuation studies in the regime $N \rightarrow \infty$, it is convenient to construct the particle system of interest as follows.

Let $\mathbf{K}_N$ be set of all bijections between $\{1, \dots, N\}$ and itself, and let its power set be $\mathcal{K}_N$. Let $\mathbf{Z}_N := (\mathbf{X}^N \times \mathbf{K}_N)^{\mathbb{N}}$ be the set of infinite sequences valued in $\mathbf{X}^N \times \mathbf{K}_N$, endow it with the σ -algebra $\mathcal{Z}_N := (\mathcal{X}^N \otimes \mathcal{K}_N)^{\otimes \mathbb{N}}$, so as to form a measurable space $(\mathbf{Z}_N, \mathcal{Z}_N)$. Let $\left\{(\widehat{\zeta}_n, \kappa_n); n \geq 0\right\}$ be the coordinate process on $\mathbf{Z}_N$

We shall write

$$\widehat{\eta}_n^N := \frac{1}{N} \sum_{i=1}^N \delta_{\widehat{\zeta}_n^i} \quad n \geq 0.$$

$$\phi_n^{\omega, N} := \frac{\widehat{\eta}_{n-1}^N Q^{\theta^{n-1}\omega}(\psi^{\theta^n\omega})}{\widehat{\eta}_{n-1}^N(G^{\theta^{n-1}\omega})} \frac{1}{\widehat{\eta}_n^N(\psi^{\theta^n\omega})} = \frac{\Phi^{\theta^{n-1}\omega}(\widehat{\eta}_{n-1}^N)(\psi^{\theta^n\omega})}{\widehat{\eta}_n^N(\psi^{\theta^n\omega})}, \quad n \geq 1.$$

$$\gamma_0^\omega := \mu_0, \quad \gamma_n^\omega(\varphi) := \eta_n^\omega(\varphi) \prod_{p=0}^{n-1} \eta_p^\omega(G^{\theta^p\omega}), \quad n \geq 1,$$

$$\widehat{\gamma}_0^{\omega, N} := \widehat{\eta}_0^N, \quad \widehat{\gamma}_n^N(\varphi) := \widehat{\eta}_n^N(\varphi) \prod_{p=0}^{n-1} \widehat{\eta}_p^N(G^{\theta^p\omega}) \phi_{p+1}^{\omega, N}, \quad n \geq 1.$$

Now for some fixed $\omega \in \Omega$, let us introduce a probability measure, $\mathbf{P}_N^\omega$, on $(\mathbf{Z}_N, \mathcal{Z}_N)$ according to the following prescription. Under $\mathbf{P}_N^\omega$, the sequences $\{\widehat{\zeta}_n; n \geq 0\}$ and $\{\kappa_n; n \geq 0\}$ are independent. The sequence $\{\widehat{\zeta}_n; n \geq 0\}$ is a Markov chain with the following law. At time $n = 0$, the random variables $\{\widehat{\zeta}_0^i\}_{i=1}^N$ are independent and identically distributed according to μ_0 . At time $n \geq 1$, the random variables $\{\widehat{\zeta}_n^i\}_{i=1}^N$ are conditionally independent given $\widehat{\zeta}_{n-1}$, and

$$\widehat{\zeta}_n^1 | \widehat{\zeta}_{n-1} \sim \widehat{\Phi}^{\theta^{n-1}\omega}(\widehat{\eta}_{n-1}^N), \quad \widehat{\zeta}_n^i | \widehat{\zeta}_{n-1} \sim \Phi^{\theta^{n-1}\omega}(\widehat{\eta}_{n-1}^N), \quad i = 2, \dots, N. \quad (47)$$

Finally, the components of the sequence $\{\kappa_n; n \geq 0\}$ are independent and identically distributed according to the uniform distribution on $(\mathbf{K}_N, \mathcal{K}_N)$. Expectation under $\mathbf{P}_N^\omega$ will be denoted by $\mathbf{E}_N^\omega$.

Remark 7. Our interest in this construction is that if we define, for each $n \geq 0$ and $i = 1, \dots, N$, the random variables

$$\zeta_n^i := \widehat{\zeta}_{\kappa_n(i)}^i$$

we obviously have the identity of empirical measures

$$\widehat{\eta}_n^N = \frac{1}{N} \sum_{i=1}^N \delta_{\zeta_n^i}, \quad n \geq 0. \quad (48)$$

Furthermore, using the fact that with $\widetilde{\mathbf{M}}$ as in (46), for any $(\omega, A) \in \Omega \times \mathcal{X}$, $\widetilde{\mathbf{M}}^\omega(x, A)$ is invariant to permutations of the coordinates $x = (x^1, \dots, x^N)$, it is straightforward to check that under $\mathbf{P}_N^\omega$, the process $\{\zeta_n; n \geq 0\}$ is Markov, with

$$\zeta_n | \zeta_{n-1} \sim \widetilde{\mathbf{M}}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot), \quad n \geq 1, \quad (49)$$

which is exactly the transition law of interest. The important thing here is that the identity of random measures (48) permits us, by construction, to perform asymptotic analysis of functionals of the empirical

measures $\left\{N^{-1} \sum_{i=1}^N \delta_{\zeta_n^i}; n \geq 0\right\}$ through study of the random variables $\left\{\widehat{\zeta}_n; n \geq 0\right\}$ rather than $\{\zeta_n; n \geq 0\}$: the conditional independence structure of the former as per (47) is easier to work with than that of the latter under (49).

To connect with the notation of previous sections, we note that under the above definitions, and for $\widetilde{\mathbf{M}}$ as in (46), we have

$$\begin{aligned}\phi_n^{\omega, N} &= \frac{\widehat{\eta}_{n-1}^N Q^{\theta^{n-1}\omega}(\psi^{\theta^n\omega})}{\widehat{\eta}_{n-1}^N(G^{\theta^{n-1}\omega})} \frac{1}{\widehat{\eta}_n^N(\psi^{\theta^n\omega})} \\ &= \left(\frac{1}{N} \sum_{i=1}^N G^{\theta^{n-1}\omega}(\zeta_{n-1}^i) \right)^{-1} \frac{\sum_{i=1}^N Q^{\theta^{n-1}\omega}(\psi^{\theta^n\omega})(\zeta_{n-1}^i)}{\sum_{i=1}^N \psi^{\theta^n\omega}(\zeta_n^i)} \\ &\equiv \phi^{\theta^n\omega}(\zeta_{n-1}, \zeta_n),\end{aligned}$$

where in the final display, ϕ^ω is as in (39). We then have

$$\prod_{p=0}^{n-1} \mathbf{G}^{\theta^n\omega}(\zeta_n) \phi^{\theta^n\omega}(\zeta_n, \zeta_{n+1}) \equiv \prod_{p=0}^{n-1} \frac{\sum_{i=1}^N Q^{\theta^{p+1}\omega}(\psi^{\theta^{p+1}\omega})(\zeta_p^i)}{\sum_{i=1}^N \psi^{\theta^{p+1}\omega}(\zeta_{p+1}^i)} \equiv \widetilde{\gamma}_n^{\omega, N}(1).$$

Introducing the Markov kernel,

$$\widetilde{M}^\omega(x, dx') \propto M^\omega(x, dx') \psi^{\theta\omega}(x'),$$

an algorithmic procedure for sampling the particle system according to (49). Here K_n and A_n are just some auxiliary random variables introduced for algorithmic convenience.

Algorithm 3 Twisted particle algorithm

For $n = 0$,

Sample $(\zeta_0^i)_{i=1}^N \stackrel{\text{iid}}{\sim} \mu_0$

For $n \geq 1$,

Sample K_n from the uniform distribution on $\{1, \dots, N\}$

Sample A_n from the distribution on $\{1, \dots, N\}$ with probabilities proportional to

$$\left\{ Q^{\theta^{n-1}\omega}(\psi^{\theta^n\omega})(\zeta_{n-1}^1), \dots, Q^{\theta^{n-1}\omega}(\psi^{\theta^n\omega})(\zeta_{n-1}^N) \right\}$$

Sample $\zeta_n^{K_n} \left| \left\{ A_n, K_n, (\zeta_{n-1}^i)_{i=1}^N \right\} \right. \sim \widetilde{M}^{\theta^n\omega}(\zeta_{n-1}^{A_n}, \cdot)$

Sample $(\zeta_n^i)_{i \neq K_n} \left| \left\{ K_n, (\zeta_{n-1}^i)_{i=1}^N \right\} \right. \stackrel{\text{iid}}{\sim} \frac{\sum_{j=1}^N G^{\theta^{n-1}\omega}(\zeta_{n-1}^j) M^{\theta^{n-1}\omega}(\zeta_{n-1}^j, \cdot)}{\sum_{j=1}^N G^{\theta^{n-1}\omega}(\zeta_{n-1}^j)}$

3.1 Analysis for $N \rightarrow \infty$

Lemma 4. Assume **(H3)**. Then for each $\omega \in \Omega$, $n \geq 1$, $\mu \in \mathcal{P}(\mathbf{X})$ and $p \geq 1$ there exist finite constants B_n^ω and $C_{n,p}^\omega$ such that for any $\varphi \in \mathcal{L}(\mathbf{X})$,

$$\left| [\Gamma_{N,n}^\omega(\mu) - \Phi_n^\omega(\mu)](\varphi) \right| \leq \|\varphi\| \frac{B_n^\omega}{N} \quad (50)$$

$$\mathbb{E}_N^\omega \left[\left| [\widehat{\eta}_n^N - \eta_n^\omega](\varphi) \right|^p \right]^{1/p} \leq \|\varphi\| \frac{C_{n,p}^\omega}{\sqrt{N}}. \quad (51)$$

From this L_p estimate it follows by a standard Borel-Cantelli argument that

Corollary 1. For each, $n \geq 0$, $\omega \in \Omega$ and $\varphi \in \mathcal{L}(\mathbf{X})$,

$$\widehat{\eta}_n^N(\varphi) - \eta_n^\omega(\varphi) \longrightarrow 0 \quad (52)$$

$$\widehat{\gamma}_n^{\omega, N}(\varphi) - \gamma_n^\omega(\varphi) \longrightarrow 0 \quad (53)$$

almost surely, as $N \rightarrow \infty$.

Lemma 5 is an extension of the CLT of Del Moral [2004, Chapter 9, Theorem 9.3.1] to the case of random test functions and the setting of the twisted particle system. It is established by an application of Jacod and Shiryaev [2002, Theorem 3.33, p. 478].

For $d \geq 1$ and $m \geq 1$ let $\varphi : (p, q, x) \in \mathbb{N} \times \{1, \dots, m\} \times \mathbf{X} \mapsto \varphi_{p,q}(x) \in \mathbb{R}^d$ be a bounded measurable function and for each $N \geq 1$ let $\beta^N := \{\beta_{p,q}^N; (p, q) \in \mathbb{N} \times \{1, \dots, m\}\}$ be a collection of random variables on $(\mathbf{Z}_N, \mathcal{Z}_N, \mathbf{P}_N)$ such that for any $p \geq 1$ and $1 \leq q \leq m$, $\beta_{p,q}^N$ is measurable w.r.t. $\sigma(\widehat{\zeta}_0, \dots, \widehat{\zeta}_{p-1})$ and $\{\beta_{0,q}^N; q \in \{1, \dots, m\}\}$ are constants. Then let the random function $f^N : (p, x) \in \mathbb{N} \times \mathbf{X} \mapsto f_p^N(x) \in \mathbb{R}^d$ be defined by

$$f_p^N(x) := \sum_{q=1}^m \beta_{p,q}^N \varphi_{p,q}(x),$$

and for $n \geq 0$ define

$$M_n^{\omega, N}(f^N) := \sum_{p=0}^n \left[\widehat{\eta}_p^N(f_p^N) - \Gamma_N^{\theta^{p-1}\omega}(\widehat{\eta}_{p-1}^N)(f_p^N) \right],$$

with the convention that $\Gamma_N^{\theta^{-1}\omega}(\widehat{\eta}_{-1}^N)(\varphi) = \mu_0$. Clearly $\{M_n^{\omega, N}(f^N); n \geq 0\}$ is an $\mathbb{R}^d$ -valued martingale w.r.t. to the natural filtration of $\{\widehat{\zeta}_n; n \geq 0\}$.

Lemma 5. Assume **(H3)** and that there exist deterministic and finite constants $(\beta_{p,q}; (p, q) \in \mathbb{N} \times \{1, \dots, m\})$ such that for each p, q ,

$$\beta_{p,q}^N \rightarrow \beta_{p,q} \quad (54)$$

in probability as $N \rightarrow \infty$. Then with

$$f_p = \sum_{q=1}^m \beta_{p,q} \varphi_{p,q},$$

for any fixed $\omega \in \Omega$, the $\mathbb{R}^d$ -valued martingale $\{\sqrt{N} M_n^{\omega, N}(f^N); n \geq 0\}$ converges in law to an $\mathbb{R}^d$ -valued Gaussian martingale $\{M_n^\omega(f); n \geq 0\}$ such that for any $1 \leq i, j \leq d$,

$$\forall n \geq 0, \quad \langle M^\omega(f^i), M^\omega(f^j) \rangle_n = \sum_{p=0}^n \eta_p^\omega [(f_p^i - \eta_p^\omega(f_p^i))(f_p^j - \eta_p^\omega(f_p^j))].$$

Now define

$$\overline{Q}^\omega(x, dx') := \frac{\overline{Q}^\omega(x, dx')}{\eta^\omega(G^\omega)}, \quad \overline{Q}_n^\omega := \frac{Q^\omega Q^{\theta\omega} \dots Q^{\theta^{n-1}\omega}}{\prod_{p=0}^{n-1} \eta_p^\omega(G^{\theta^p\omega})},$$

and notice that

$$\mu_0 \overline{Q}_n^\omega = \eta_n^\omega.$$

Theorem 2. Assume **(H3)**. Then for any $n \geq 0$, $\omega \in \Omega$ and $\varphi \in \mathcal{L}(\mathbf{X})$,

$$\sqrt{N} [\widehat{\eta}_n^N(\varphi) - \eta_n^\omega(\varphi)] \Rightarrow \mathcal{N}(0, \sigma_{n,\omega}^2(\varphi)), \quad (55)$$

$$\sqrt{N} [\widehat{\gamma}_n^{\omega, N}(\varphi) - \gamma_n^\omega(\varphi)] \Rightarrow \mathcal{N}(0, \varsigma_{n,\omega}^2(\varphi)), \quad (56)$$

as $N \rightarrow \infty$ where

$$\sigma_{n,\omega}^2(\varphi) = \sum_{p=0}^n \eta_p^\omega \left[\left(\overline{Q}_{n-p}^{\theta^p \omega}(\varphi - \eta_n^\omega(\varphi)) \right)^2 \right]. \quad (57)$$

and

$$\varsigma_{n,\omega}^2(\varphi) = \sum_{p=0}^n \gamma_p^\omega(1)^2 \eta_p^\omega \left[\left(Q_{n-p}^{\theta^p \omega}(\varphi) - \frac{\psi^{\theta^p \omega}}{\eta_p^\omega(\psi^{\theta^p \omega})} \eta_p^\omega Q_{n-p}^{\theta^p \omega}(\varphi) \right)^2 \right]. \quad (58)$$

Remark 8. The asymptotic variance expression (57) is independent of the particular choice of ψ (subject to **(H3)** of course), and is thus exactly the same expression obtained in the CLT for the standard particles system (i.e. ψ constant) see for example [Del Moral, 2004, Proposition 9.4.2]. However, the asymptotic variance in (58) clearly does depend on ψ in general.

3.2 Analysis for $n \rightarrow \infty$

For $\widetilde{\mathbf{M}}$ as in (46), we obtain an estimate of $\Upsilon_N(\widetilde{\mathbf{M}})$ which exhibits its dependence on N and the discrepancy between ψ and h . The proof is given in section 5.

Proposition 5. Assume **(H1)**, **(H2)** and

$$\sup_{\omega, \omega', x, x'} \frac{\psi^\omega(x)}{\psi^{\omega'}(x')} < \infty.$$

Then for any $N \geq 2$,

$$\Upsilon_N(\widetilde{\mathbf{M}}) \leq \log \left[1 + \frac{1}{N-1} \mathcal{D}_{\mathbb{P}}(\psi, h) \right],$$

where

$$\mathcal{D}_{\mathbb{P}}(\psi, h) := \mathbb{P} - \text{ess sup}_\omega \left\{ \left[2 \sup_{(z, z') \in \mathbf{X}^2} \frac{\psi^\omega(z)}{\psi^\omega(z')} - 1 \right] \sup_{z \in \mathbf{X}} \left(\frac{\psi^\omega(z)}{h^\omega(z)} \right) \sup_{(z, z') \in \mathbf{X}^2} \left| \frac{h^\omega(z)}{\psi^\omega(z)} - \frac{h^\omega(z')}{\psi^\omega(z')} \right| \right\}.$$

4 Discussion

4.1 Sequential Importance Sampling

Notice that in the case $N = 1$, we have the identity $\mathbf{M}^\omega = M^\omega$. This observation leads us to consider how our results can be applied to analyze the variance growth behaviour of Sequential Importance Sampling (SIS), in the following sense.

Let $\widetilde{M} : \Omega \times \mathbf{X} \times \mathcal{X} \rightarrow [0, 1]$ be a Markov kernel, and for some $L \geq 1$ and any fixed $\omega \in \Omega$, let $\{X_n^i; n \geq 0\}_{i=1}^L$ be L iid time-inhomogeneous Markov chains, each with law

$$X_0^i \sim \mu_0, \quad X_n^i | X_{n-1}^i = x_{n-1}^i \sim \widetilde{M}^{\theta^{n-1} \omega}(x_{n-1}^i, \cdot) \quad n \geq 1.$$

Now to connect with the setting of sections 2.3 and 2.4, continue with $N = 1$, and take $\widetilde{\mathbf{M}} := \widetilde{M}$. We shall assume that **(H1)** and **(H2)** hold and that $\widetilde{M}$ is a member of $\mathbb{M}$. The quantity

$$\frac{1}{L} \sum_{i=1}^L \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p^i) \phi^{\theta^p \omega}(X_p^i, X_{p+1}^i) \right] \quad (59)$$

is clearly an unbiased estimator of $\mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p^i) \right]$. Furthermore, since the L Markov chains are independent, for any fixed ω , the relative variance of (59) is

$$\frac{1}{L} [\mathcal{V}_{n,1}^\omega + 1], \quad (60)$$

where $\mathcal{V}_{n,1}^\omega$ is as in (41). By application of Theorem 1, we find that except when (up to the sets of measure zero mentioned in Theorem 1) $\widetilde{M}^\omega(x, dx') \propto M(x, dx')h^{\theta\omega}(x')$, we have the $\mathbb{P}$ -almost-sure convergence:

$$\frac{1}{n} \log \mathcal{V}_{n,1}^\omega \rightarrow \Upsilon_1(\widetilde{M}) > 0, \quad (61)$$

and supposing that now $L = L(n)$, the leading term in (60) $\mathbb{P}$ -almost-surely behaves according to:

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \log \left(\frac{1}{L} \mathcal{V}_{n,1}^\omega \right) = \Upsilon_1(\widetilde{M}) - \limsup_{n \rightarrow \infty} \frac{1}{n} \log L(n). \quad (62)$$

In summary, whenever $\Upsilon_1(\widetilde{M}) > 0$, it is necessary that $L(n)$ must be scaled up exponentially in n in order to prevent exponential growth of the relative variance of (59). In this sense the SIS approach is typically an inefficient method for approximating $\mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p \omega}(X_p^i) \right]$, at least relative to particle methods, which we shall now discuss.

4.2 The bootstrap particle filter

In the case

$$G(\omega, x) := g(x, Y_0(\omega)), \quad M(\omega, x, dx') := f(x, dx'), \quad (63)$$

we have that $\{\mathbf{M}^\omega; \omega \in \Omega\}$ is the collection of the transitions of the bootstrap particle filter, as described in the introduction. When **(H1)** and **(H2)** hold, by Lemma 3 we find that in this scenario, for any $N \geq 1$,

$$\Upsilon_N(\mathbf{M}) = 0 \quad \text{if and only if for } \mathbb{P} - a.a. \omega, \exists A_\omega \in \mathcal{X} \text{ s.t. } \nu(A_\omega^c) = 0 \text{ and } g(x, Y_0(\omega)) \text{ is constant on } A_\omega.$$

The condition of $g(x, Y_0(\omega))$ being constant in x represents an entirely degenerate HMM in which the observations do not provide any information about the hidden state. Thus we concentrate on the situation $\Upsilon_N(\mathbf{M}) > 0$. By an application of Proposition 5 in the case that $\psi^\omega(x) = 1$ for all ω and x , and using the bound (30) of Proposition 2 we find that there exists a constant $c < \infty$ such that

$$\frac{1}{n} \log \mathcal{V}_{n,N}^\omega \rightarrow \Upsilon_N(\mathbf{M}) \leq \log \left[1 + \frac{c}{N-1} \right], \quad (64)$$

where the convergence holds $\mathbb{P}$ -almost surely. The practical importance of this result is that it shows why even the rather basic bootstrap filter is to be preferred over the SIS method in terms of variance-growth behaviour, as seen by comparing (64) with (62).

It should be noted that under our assumptions **(H1)** and **(H2)**, the bound (64) is implied by the bounds of Cérou et al. [2011, Theorem 5.1]. The results of Cérou et al. [2011] provide important information about the non-asymptotic-in- n behaviour of the relative variance, which our Proposition 5 does not. On the other hand, Proposition 5 applies not just to the standard particle transition $\mathbf{M}$, but also to twisted transitions, to which the analysis of Cérou et al. [2011] does not extend. We shall now discuss these twisted transitions.

Continuing with the setting (63), and assuming that **(H1)** and **(H2)** hold, we shall now provide some interpretation of h . The objects appearing in part 1) of Proposition 2 have the following interpretations: $\Phi_n^{\theta^{-n}\omega}(\mu)$ is the one-step-ahead prediction filter initialized at time $-n$ using μ , and run forward to time zero, thus conditioning on the observations $Y_{-n}(\omega), \dots, Y_{-1}(\omega)$. Let us write this prediction filter probability measure as $P_n^\omega \equiv \Phi_n^{\theta^{-n}\omega}(\mu)$. The quantity $Q_n^\omega(1)(x)$ is the conditional likelihood of observations $Y_0(\omega), \dots, Y_{n-1}(\omega)$ given that the hidden state in the HMM at time zero is x . Thus if we denote by Π_n^ω the probability measure given by

$$\Pi_n^\omega(A) := \frac{\int_A P_n^\omega(dx) Q_n^\omega(1)(x)}{\int_{\mathcal{X}} P_n^\omega(dz) Q_n^\omega(1)(z)}, \quad A \in \mathcal{X},$$

we find that by inspection of part 1) of Proposition 2, that h can be interpreted as the point-wise limit

$$h(\omega, x) \equiv \lim_{n \rightarrow \infty} \frac{d\Pi_n^\omega}{dP_n^\omega}(x). \quad (65)$$

Moreover, by part 2) of Proposition 2 we find that

$$\sup_{\omega, x} \left| \frac{d\Pi_n^\omega}{dP_n^\omega}(x) - h(\omega, x) \right| \leq C\rho^n \quad (66)$$

for some constants $C < \infty$ and $\rho \in (0, 1)$.

Let us now consider a twisted bootstrap particle filter (as per the Section 1), in the case that for some fixed $\ell \geq 1$, we take $\psi^\omega := d\Pi_\ell^\omega/dP_\ell^\omega$, and as an instance of the setup in section 3, we let $\widetilde{\mathbf{M}}$ be as per (46) with this choice of ψ^ω . Although typically unavailable in practice, this ψ^ω allows an illustrative application of Proposition 5. Indeed, using (66), and the fact that under the bounds of part 3) of Proposition 2 $h(\omega, x)$ is uniformly bounded above and below away from zero, elementary manipulations show that there exists some finite constant $C' < \infty$ such that

$$\Upsilon_N(\widetilde{\mathbf{M}}) \leq \log \left[1 + \frac{C'\rho^\ell}{N-1} \right]. \quad (67)$$

We see that, in principle, increasing the lag length ℓ is useful in helping to control $\Upsilon_N(\widetilde{\mathbf{M}})$.

Now under the mild regularity condition **(H3)**, for fixed ω and n , and φ a bounded measurable function on $\mathbf{X}$, Corollary 1 shows that for the twisted particle filter,

$$N^{-1} \sum_{i=1}^N \varphi(\zeta_n^i) - \pi_n^\omega(\varphi) \rightarrow 0 \quad (68)$$

as $N \rightarrow \infty$, with probability one, independently of ψ . Furthermore, by Theorem 2, $N^{-1/2} \sum_{i=1}^N [\varphi(\zeta_n^i) - \pi_n^\omega(\varphi)]$ converges in distribution to a centered Gaussian random variable with variance independent of ψ , i.e. the same asymptotic variance obtained under the standard bootstrap particle filter.

4.3 Auxiliary particle filters

In addition to the ingredients of the HMM given in section 1, introduce $r : \Omega \times \mathbf{X} \rightarrow \mathbb{R}_+$ such that for each ω , $r^\omega(x)$ is strictly positive and bounded in x . Then set:

$$G^\omega(x) := \frac{g(x, Y_0(\omega)) \int_{\mathbf{X}} r^{\theta\omega}(z) f(x, dz)}{r^\omega(x)}, \quad M^\omega(x, dx') \propto f(x, dx') r^{\theta\omega}(x'). \quad (69)$$

In this case, sampling according to the corresponding $\mathbf{M}^\omega$, we obtain a form of Auxiliary Particle Filter (APF) [Pitt and Shephard, 1999, Johansen and Doucet, 2008, Douc et al., 2009]. More specifically, let $\{\mu_0^\omega \in \mathcal{P}(\mathbf{X}); \omega \in \Omega\}$ be the family of probability measures such that $\mu_0^\omega(dx) \propto r^\omega(x) \mu_0(dx)$, where μ_0 is the initial distribution in the HMM, as in section 1. Then sampling

$$\zeta_0 \sim (\mu_0^\omega)^{\otimes N}, \quad \zeta_n | \zeta_{n-1} \sim \mathbf{M}^{\theta^{n-1}\omega}(\zeta_{n-1}, \cdot), \quad (70)$$

it is straightforward to check that

$$\frac{1}{\mu_0(r^\omega)} \prod_{p=0}^{n-1} \mathbf{G}^{\theta^p\omega}(\zeta_p) \left(\frac{1}{N} \sum_{i=1}^N \frac{1}{r^{\theta^n\omega}(\zeta_n^i)} \right) \quad (71)$$

is an unbiased estimator of Z_n^ω , at least whenever the expectation of (71) is finite. If, for example, r is bounded below away from zero, and **(H1)** and **(H2)** hold, then Proposition 4 may be applied to establish the existence of $\Upsilon_N(\mathbf{M}) \geq 0$ such that the following convergence holds $\mathbb{P}$ -almost surely:

$$\frac{1}{n} \log \mathbb{E}_N^\omega \left[\frac{1}{\mu_0(r^\omega)^2} \left(\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p\omega}(\zeta_p) \right)^2 \left(\frac{1}{N} \sum_{i=1}^N \frac{1}{r^{\theta^n\omega}(\zeta_n^i)} \right)^2 \right] - \frac{2}{n} \log Z_n^\omega \rightarrow \Upsilon_N(\mathbf{M}),$$

since the $\mu_0(r^\omega)$ and $N^{-1} \sum_{i=1}^N [r^{\theta^n\omega}(\zeta_n^i)]^{-1}$ terms have no asymptotic contribution and since the convergence in Proposition 4 is independent of the distribution from which the particle system is initialized. If

we choose $r(\omega, x) := g(x, Y_0(\omega))$ we obtain the “fully adapted” APF of Pitt and Shephard [1999]. Lemma 3 then shows that $\Upsilon_N(\mathbf{M}) = 0$ if and only if $\mathbb{P}$ -almost all ω , $\int_{\mathbf{X}} g(z, Y_1(\omega)) f(x, dz)$ is ν -almost everywhere a constant. The latter condition usually does not hold in situations of practical interest. One may follow the generic recipe of section 3 and derive a twisted version of this full-adapted APF, but let us consider a different choice of r . Under the now familiar regularity conditions, the Ω -indexed family of kernels given by

$$g(x, Y_0(\omega)) f(x, dx') \quad (72)$$

have an associated eigen-function given by the point-wise limit on the right hand side of (65). Let us then choose

$$r(\omega, x) := \lim_{n \rightarrow \infty} \frac{d\Pi_n^\omega}{dP_n^\omega}(x), \quad (73)$$

so that

$$\int_{\mathbf{X}} g(x, Y_0(\omega)) f(x, dx') r^{\theta\omega}(x') = \chi_\omega r^\omega(x)$$

for a non-negative random variable χ which is the eigen-value for the family of kernels (72). In this case we find that the G^ω given in (69) is constant in x and Lemma 3 then shows that $\Upsilon_N(\mathbf{M}) = 0$, and the eigen-function of Q^ω is essentially constant. The interpretation of this result is that the variance growth rate behaviour of (71), with the particle system sampled according to the APF with r given by (73), cannot be improved upon by twisting.

Now let us consider the empirical measures obtained under the APF, for some general choice of r . If one wishes to use $N^{-1} \sum_{i=1}^N \delta_{\zeta_n^i}$ to approximate the prediction filter π_n^ω in a $N \rightarrow \infty$ consistent manner then, in general and in contrast to (68), some re-weighting must be applied. For example, assuming **(H3)** holds with G as in (69), Corollary 1 and some elementary manipulations involving the Φ_n^ω semi-group show that for bounded measurable φ ,

$$\frac{\sum_{i=1}^N \varphi(\zeta_n^i) / r^{\theta^n \omega}(\zeta_n^i)}{\sum_{i=1}^N 1 / r^{\theta^n \omega}(\zeta_n^i)} - \pi_n^\omega(\varphi) \rightarrow 0, \quad (74)$$

as $N \rightarrow \infty$, with probability 1 under the law of the particle system specified by (69)-(70). Of course, (74) can be verified using the results of Johansen and Doucet [2008], Douc et al. [2009]. However, the new perspective which our results bring here is that, as we have seen in (67) and (68), the twisted bootstrap particle filter can in theory achieve $\Upsilon_N(\widetilde{\mathbf{M}}) = 0$ whilst not requiring re-weighting of particles to obtain consistent approximations of π_n^ω .

4.4 Numerical illustrations

In order to give some impression of the practical performance of the algorithms we have analyzed, we now present some numerical findings. **(H2)** is not satisfied for the M and G which specify the models below; in this section some of our theoretical findings can only be used as guidelines for the design of practical algorithms. We note however, that the much milder regularity condition **(H3)** is satisfied for the models we consider and thus Corollary 1 and Theorem 2 apply to the particle systems in question.

We shall first consider the influence of ψ on the variance-growth behaviour of the twisted bootstrap particle filter (henceforth TPF). The purpose of this example is to illustrate an idealized scenario in which $\psi^\omega := d\Pi_\ell^\omega / dP_\ell^\omega$ can be computed exactly. Consider a linear-Gaussian state-space model where $X_n = 0.9X_{n-1} + V_n$, $Y_n = X_n + W_n$ where $(V_n), (W_n)$ are iid zero mean Gaussian sequences. We shall consider a TPF with $\psi^\omega = d\Pi_\ell^\omega / dP_\ell^\omega$ for some $\ell \geq 0$, which can be computed in closed form for this model. Note that the TPF algorithm can be implemented with ψ^ω only known up to a constant of proportionality. Figure 1 shows variance growth behaviour estimated empirically using 2000 independent runs of the algorithm for a single observation sequence, which was drawn from the model and then fixed. Convergence of $n^{-1} \log \widetilde{\mathcal{V}}_{n,N}^\omega$ is apparent and the influence of ℓ on the rate of variance growth is substantial.

We now turn to a standard stochastic volatility model, in which $d\Pi_\ell^\omega / dP_\ell^\omega$ is unavailable in closed form, but for which a standard deterministic (henceforth: “the”) approximation is available. For details of the model, this approximation and a real data of daily returns on pound/dollar exchange rates see [Doucet et al., 2006] and references therein. We tested the TPF and APF using this data set and the same model




Figure 1: Linear-Gaussian model and the TPF. Left: estimated values of $\tilde{\mathcal{V}}_{n,N}^\omega$ against n . Right: estimated values of $n^{-1} \log \tilde{\mathcal{V}}_{n,N}^\omega$ against n . In both cases $N = 100$ and ω is fixed. Lag-lengths: red, $\ell = 0$ (i.e. the bootstrap filter); yellow, $\ell = 1$; green, $\ell = 2$; cyan, $\ell = 5$. The $\ell = 0$ plot is omitted from the right-hand figure due to scale constraints. To the precision of these figures, increasing the lag beyond $\ell = 5$ had no noticeable influence on the variance.




Figure 2: Stochastic Volatility. Estimated values of $\tilde{\mathcal{V}}_{n,N}^\omega$ against n , with $N = 1000$ and ω fixed. Left: TPF. Right: APF. Red, $\ell = 0$ (the bootstrap algorithm), yellow, $\ell = 1$, cyan, $\ell = 2$, blue, $\ell = 5$; green, $\ell = 10$; violet, $\ell = 50$.




Figure 3: Stochastic Volatility model. Left: estimated $\tilde{\mathcal{V}}_{n,N}^\omega$ for the TPF against n for $\ell = 5$, ω fixed and: red, $N = 10$; yellow, $N = 100$; green $N = 1000$. Right: empirical variance of particle approximations of the mean of π_n^ω against n , with $N = 1000$ and ω fixed. Cyan, TPF; red, APF. For both algorithms, $\ell = 5$.

parameter settings as in the aforementioned paper. We took both ψ^ω (for the twisted bootstrap filter) and r^ω (for the APF as in (69)) to both be the approximation of $d\Pi_\ell^\omega/dP_\ell^\omega$.

Figure 2 shows empirical variance-growth behaviour for a range of values of ℓ , estimated from 2000 independent runs of each algorithm. For both algorithms, increasing ℓ appears to generally yield a decrease in variance. The figures indicate that apart from occasional fluctuations, the APF mostly exhibits lower variance than the TPF, however, we found this phenomenon to be dependent on model parameter settings, for other parameter values we found the TPF exhibited lower variance than the APF (not shown).

The left plot in Figure 3 illustrates how the variance of the estimates from the TPF varies with N . An increase in variance growth rate is evident as N is decreased. The right plot of Figure 3 shows the empirical variance of particle estimates of the mean of the prediction filter π_n^ω against n , obtained from the TPF and APF both with $\ell = 5$. It is notable that here the TPF generally exhibits lower variance than the APF. Results for the standard bootstrap particle filter were found to be identical to those for the TPF on the scale of this figure, this is in agreement with the conclusions of Theorem 2 applied to the TPF - that the asymptotic variance of prediction filter approximations is independent of ψ .

4.5 Generalizations and extensions

- We have only mentioned the multinomial resampling scheme, appearing implicitly in the definition of $\mathbf{M}$ given in (33). Several alternative schemes are popular in practice. In order to develop extensions of Proposition 4 and Theorem 1 to alternative schemes (assuming resampling is applied at every time step, and with a fixed number of particles) it suffices to re-define $\mathbf{M}$ appropriately so that it incorporates the resampling scheme of interest, check that the conditions in the statement of Lemma 1 are satisfied, and check that Lemma 2 holds with $\mathbf{Q}$ re-defined in terms of this new $\mathbf{M}$. Of course, the new $\mathbf{M}$ will influence the form of the corresponding twisted algorithms.
- Some types of standard particle algorithm, and variants of the APF, resample according to weights which depend on two or more historical components of the trajectory of each particle. Such algorithms can be incorporated into the framework presented here by a simple state-space augmentation. For example, starting from each Markov kernel M^ω on $(\mathbf{X}, \mathcal{X})$ (according to which particles are sampled in the algorithm of interest) one builds a kernel, $\bar{M}^\omega(x, dz) := \delta_{x_2}(dz_1)M^\omega(z_1, dz_2)$ on $(\mathbf{X}^2, \mathcal{X}^{\otimes 2})$, where

$x = (x_1, x_2), z = (z_1, z_2)$ are points in X^2 , and introduces the appropriate incremental weight $\overline{G}^\omega(x)$. Then the analyses of Section 2 can be repeated with only superficial differences: when M^ω satisfies (18), then $\overline{M}^\omega$ satisfies a 2-step version of the same condition, one simply then works on $(X^2, \mathcal{X}^{\otimes 2})$ instead of $(X, \mathcal{X})$, dealing with the kernel $\overline{Q}^\omega(x, dz) := \overline{G}^\omega(x) \overline{M}^\omega(x, dz)$ instead of Q^ω .

5 Proofs and auxiliary results

Proofs for section 2.2

Before presenting the proof of Proposition 2 it is opportune to observe that for any $n \geq 1$:

$$\Phi_n^\omega = \Phi_{n-1}^{\theta\omega} \circ \Phi^\omega. \quad (75)$$

The formula is validated by noticing that $\Phi_0^\omega = Id$, $\Phi_1^\omega = \Phi^\omega$, and when (75) holds at rank n , using the definition of Φ_{n+1}^ω , composing $\Phi^{\theta^n\omega}$ on the left of the objects appearing in (75) and then using the definition of $\Phi_n^{\theta\omega}$ gives the equalities:

$$\Phi_{n+1}^\omega = \Phi^{\theta^n\omega} \circ \Phi_n^\omega = \Phi^{\theta^n\omega} \circ \Phi_{n-1}^{\theta\omega} \circ \Phi^\omega = \Phi_n^{\theta\omega} \circ \Phi^\omega,$$

so that the formula (75) holds at rank $n+1$. A simple inductive arguments then shows that for any $n, m \geq 1$ and any $\omega \in \Omega$,

$$\Phi_{n+m}^\omega = \Phi_n^{\theta^m\omega} \circ \Phi_m^\omega. \quad (76)$$

Proof. (of Proposition 2). Throughout the proof C is a finite constant whose value may change on each appearance.

We first address (26) and (28). Applying (76) with $\theta^{-n-m}\omega$ in place of ω gives

$$\Phi_{n+m}^{\theta^{-n-m}\omega} = \Phi_n^{\theta^{-n}\omega} \circ \Phi_m^{\theta^{-n-m}\omega},$$

so for any $\mu \in \mathcal{P}(X)$, taking $\mu' = \Phi_m^{\theta^{-n-m}\omega}(\mu)$ then applying (25) we obtain, for any $n, m \geq 1$,

$$\sup_{\omega \in \Omega} \sup_{\mu \in \mathcal{P}(X)} \left| \left[\Phi_n^{\theta^{-n}\omega}(\mu) - \Phi_{n+m}^{\theta^{-n-m}\omega}(\mu) \right] (\varphi) \right| \leq \|\varphi\| C \rho^n. \quad (77)$$

Taking $\varphi = \mathbb{I}_A$ for any $A \in \mathcal{X}$, (77) shows that for some fixed μ and ω , $\left\{ \Phi_n^{\theta^{-n}\omega}(\mu)(A); n \geq 0 \right\}$ is a real-valued Cauchy sequence and together with [Grey, 2001, Theorem 1] this establishes the existence of $\eta^\omega \in \mathcal{P}(X)$ such that (26) holds. Moreover, taking $m \rightarrow \infty$ in (77) we then find that

$$\sup_{\omega \in \Omega} \sup_{\mu \in \mathcal{P}(X)} \left| \left[\Phi_n^{\theta^{-n}\omega}(\mu) - \eta^\omega \right] (\varphi) \right| \leq \|\varphi\| C \rho^n, \quad (78)$$

and thus $\Phi_n^{\theta^{-n}\omega}(\mu)$ converges in total variation to η^ω , uniformly over μ and ω . This establishes (28).

We next address (27) and (29). We shall establish that

$$\sup_{(\omega, x) \in \Omega \times X} \sup_{\mu \in \mathcal{P}(X)} \left| \frac{Q_{n+1}^\omega(1)(x)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu) Q_{n+1}^\omega(1)} - \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\mu) Q_n^\omega(1)} \right| \leq C \rho^n. \quad (79)$$

First note that under **(H2)**,

$$\tilde{h}_{n,\mu}^\omega(x) := \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\mu) Q_n^\omega(1)}$$

satisfies for any $\omega, \omega', x, x', \mu$,

$$\sup_{n \geq 1} \frac{\tilde{h}_{n,\mu}^\omega(x)}{\tilde{h}_{n,\mu}^{\omega'}(x')} = \sup_{n \geq 1} \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\mu) Q_n^\omega(1)} \frac{\Phi_n^{\theta^{-n}\omega'}(\mu) Q_n^{\omega'}(1)}{Q_n^{\omega'}(1)(x')} \leq \left(\beta \frac{\epsilon_+}{\epsilon_-} \right)^2. \quad (80)$$

We have

$$\begin{aligned}
& \left| \frac{Q_{n+1}^\omega(1)(x)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)} - \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)} \right| \\
&= \left| \frac{Q_{n+1}^\omega(1)(x)\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1) - Q_n^\omega(1)(x)\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)} \right| \\
&\leq \left| \frac{Q_{n+1}^\omega(1)(x)\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1) - Q_n^\omega(1)(x)\Phi_n^{\theta^{-n}\omega}(\mu)Q_{n+1}^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)} \right| \\
&\quad + \left| \frac{Q_n^\omega(1)(x)\Phi_n^{\theta^{-n}\omega}(\mu)Q_{n+1}^\omega(1) - Q_n^\omega(1)(x)\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)} \right| \\
&= \frac{Q_n^\omega(1)(x)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)} \left| \frac{Q_{n+1}^\omega(1)(x)\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1) - Q_n^\omega(1)(x)\Phi_n^{\theta^{-n}\omega}(\mu)Q_{n+1}^\omega(1)}{Q_n^\omega(1)(x)\Phi_n^{\theta^{-n}\omega}(\mu)Q_{n+1}^\omega(1)} \right| \\
&\quad + \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)} \left| \frac{\Phi_n^{\theta^{-n}\omega}(\mu)Q_{n+1}^\omega(1) - \Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)} \right| \\
&= \tilde{h}_{n,\mu}^\omega(x) \frac{\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)} \left| \frac{Q_{n+1}^\omega(1)(x)}{Q_n^\omega(1)(x)} - \frac{\Phi_n^{\theta^{-n}\omega}(\mu)Q_{n+1}^\omega(1)}{\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)} \right| \\
&\quad + \tilde{h}_{n,\mu}^\omega(x) \left| \frac{\Phi_n^{\theta^{-n}\omega}(\mu)Q_{n+1}^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)} - \frac{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)} \right| \\
&= \tilde{h}_{n,\mu}^\omega(x) \frac{\Phi_n^{\theta^{-n}\omega}(\mu)Q_n^\omega(1)}{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)Q_{n+1}^\omega(1)} \left| \left[\Phi_n^\omega(\delta_x) - \left(\Phi_n^\omega \circ \Phi_n^{\theta^{-n}\omega} \right) (\mu) \right] Q^{\theta^n\omega}(1) \right| \\
&\quad + \tilde{h}_{n,\mu}^\omega(x) \left| \left[\Phi_n^{\theta^{-n}\omega}(\mu) - \left(\Phi_n^{\theta^{-n}\omega} \circ \Phi_n^{\theta^{-n-1}\omega} \right) (\mu) \right] (\tilde{h}_{n+1,\mu}^\omega) \right| \\
&\leq C\rho^n \quad \forall x, \omega, \mu,
\end{aligned}$$

where (80), **(H2)** and (25) have been applied. Thus (79) is proved and then for any $m, n \geq 1$,

$$\sup_{\omega, x, \mu} \left| \tilde{h}_{n,\mu}^\omega(x) - \tilde{h}_{n+m,\mu}^\omega(x) \right| \leq C \sum_{q=0}^{m-1} \rho^{n+q} \leq \rho^n \frac{C}{1-\rho}.$$

Then $\{\tilde{h}_{n,\mu}^\omega(x); n \geq 1\}$ is Cauchy and real-valued, which is enough to establish the existence of $h : \Omega \times \mathbf{X} \rightarrow \mathbb{R}_+$ such that (27) holds, and furthermore

$$\sup_{\omega, x, \mu} \left| \tilde{h}_{n,\mu}^\omega(x) - h^\omega(x) \right| \leq C\rho^n, \tag{81}$$

which establishes (29). The measurability of h stated in part 1) holds as it is the point-wise limit of a sequence of $\mathcal{F} \otimes \mathcal{X}$ -measurable functions.

Turning now to prove part 3) of the Proposition, note that by (78), $\lambda_\omega = \eta_\omega(G^\omega) = \lim_{n \rightarrow \infty} \Phi_n^{\theta^{-n}\omega}(\mu)(G^\omega)$, i.e. λ is the point-wise limit of a sequence of measurable functions, and is therefore measurable. The λ part of (30) holds immediately under **(H2)**, and the h part of (30) holds due to (80) and (81).

We now turn to the proof of part 4). We now establish that the triple (η, h, λ) does indeed satisfy (31). Firstly, for the measure equation, we have

$$\frac{\Phi_n^{\theta^{-n}\omega}(\mu)Q^\omega(A)}{\Phi_n^{\theta^{-n}\omega}(\mu)(G^\omega)} = \Phi^\omega(\Phi_n^{\theta^{-n}\omega}(\mu))(A) = \Phi_{n+1}^{\theta^{-n-1}\omega}(\mu)(A). \tag{82}$$

By the strong convergence in (78), and the fact that under **(H2)**, $\sup_x Q^\omega(A)(x) \leq \sup_x G^\omega(x) < \infty$, the left hand side of (82) converges to $\eta^\omega Q^\omega(A)/\lambda_\omega$ and the right hand side of (82) converges to $\eta^{\theta^\omega}(A)$. Thus η

satisfies the first equation in (31). For the second equation in (31), choose $n \geq 1$ arbitrarily and notice that

$$Q^\omega \left(\tilde{h}_{n,\mu}^{\theta\omega} \right) (x) = \frac{Q^\omega Q_n^{\theta\omega}(1)(x)}{\Phi_n^{\theta^{-n+1}\omega}(\mu) Q_n^{\theta\omega}(1)} \quad (83)$$

$$\begin{aligned} &= \frac{Q_{n+1}^\omega(1)(x)}{\Phi_n^{\theta^{-n+1}\omega}(\mu) Q_n^{\theta\omega}(1)} \\ &= \tilde{h}_{n+1,\mu}^\omega(x) \frac{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu) Q_{n+1}^\omega(1)}{\Phi_n^{\theta^{-n+1}\omega}(\mu) Q_n^{\theta\omega}(1)} \\ &= \tilde{h}_{n+1,\mu}^\omega(x) \frac{\Phi_{n+1}^{\theta^{-n-1}\omega}(\mu) Q_{n+1}^\omega(1)}{\Phi_{n-1}^{\theta^{-n+1}\omega}(\mu) Q_n^{\theta\omega}(1)} \cdot \Phi_{n-1}^{\theta^{-n+1}\omega}(\mu) Q^\omega(1) \\ &= \tilde{h}_{n+1,\mu}^\omega(x) \frac{\Phi_{n-1}^{\theta^{-n+1}\omega} \left[\Phi_2^{\theta^{-n-1}\omega}(\mu) \right] Q_{n+1}^\omega(1)}{\Phi_{n-1}^{\theta^{-n+1}\omega}(\mu) Q_{n+1}^\omega(1)} \cdot \Phi_{n-1}^{\theta^{-n+1}\omega}(\mu) Q^\omega(1) \\ &\rightarrow h^\omega(x) \cdot 1 \cdot \lambda_\omega, \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (84)$$

where the convergence in (84) is due to (81); (25) applied in conjunction with the property

$$\sup_{n \geq 1} \sup_{x, x'} \frac{Q_{n+1}^\omega(1)(x)}{Q_{n+1}^\omega(1)(x')} < \infty$$

which holds under **(H2)**; and (28). Furthermore the l.h.s. of (83) converges to $Q^\omega(h^{\theta\omega})(x)$ by (81), (80) and the dominated convergence theorem. This verifies that the second equation of (31) is satisfied. For the third equation, we have, for any $n \geq 1$,

$$\begin{aligned} |\eta^\omega(h^\omega) - 1| &= \left| \eta^\omega(h^\omega) - \Phi_n^{\theta^{-n}\omega}(\mu) (\tilde{h}_{n,\mu}^\omega) \right| \\ &\leq \left| \left[\eta^\omega - \Phi_n^{\theta^{-n}\omega}(\mu) \right] (h^\omega) \right| + \sup_x \left| h^\omega(x) - \tilde{h}_{n,\mu}^\omega(x) \right| \end{aligned}$$

and since we have already proved part 3), which implies $\sup_x h^\omega(x) < \infty$, the convergence in (78) and (81) show it must be the case that $\eta^\omega(h^\omega) = 1$.

Now for the uniqueness element of part 4). Suppose that there exists a triple $(\bar{\eta}, \bar{h}, \bar{\lambda})$ of the desired nature and such that

$$\bar{\eta}^\omega Q^\omega = \bar{\lambda}_\omega \bar{\eta}^{\theta\omega}, \quad Q^\omega(\bar{h}^{\theta\omega}) = \bar{\lambda}_\omega \bar{h}^\omega, \quad \bar{\eta}^\omega(\bar{h}^\omega) = 1, \quad \text{for all } \omega \in \Omega. \quad (85)$$

Then integrating the first equation in (85), we have $\bar{\lambda}_\omega = \bar{\eta}_\omega(G^\omega)$, because $\bar{\eta}_\omega$ is, by hypothesis, a probability measure. Thus $\Phi^\omega(\bar{\eta}^\omega) = \bar{\eta}^{\theta\omega}$ for any ω , so via iteration we find $\Phi_n^{\theta^{-n}\omega}(\bar{\eta}^{\theta^{-n}\omega}) = \bar{\eta}^\omega$, and strong convergence of (78) then demands that $\bar{\eta}^\omega = \eta^\omega$. Thus $\bar{\eta} = \eta$ and therefore $\bar{\lambda} = \lambda$. It remains to show that $\bar{h} = h$. To this end, first note that

$$\begin{aligned} \left| \frac{Q_n^\omega(\varphi)(x)}{\prod_{p=0}^{n-1} \lambda_{\theta^p\omega}} - h^\omega(x) \eta^{\theta^n\omega}(\varphi) \right| &\leq \left| \frac{Q_n^\omega(\varphi)(x)}{Q_n^\omega(1)(x)} - \eta^{\theta^n\omega}(\varphi) \right| \beta \frac{\epsilon_+}{\epsilon_-} \\ &\quad + \|\varphi\| \left| \frac{Q_n^\omega(1)(x)}{\Phi_n^{\theta^{-n}\omega}(\eta^{\theta^{-n}\omega}) Q_n^\omega(1)} - h^\omega(x) \right| \\ &\rightarrow 0, \quad \text{as } n \rightarrow \infty \end{aligned} \quad (86)$$

where the identity $\prod_{p=0}^{n-1} \lambda_{\theta^p\omega} = \eta^\omega Q_n^\omega(1) = \Phi_n^{\theta^{-n}\omega}(\eta^{\theta^{-n}\omega}) Q_n^\omega(1)$, which holds due to the measure equation in (31); then **(H2)**; and then (78) and (81) have been applied. But, under the hypotheses that $Q^\omega(\bar{h}^{\theta\omega}) = \bar{\lambda}_\omega \bar{h}^\omega$ and $\bar{\eta}^\omega(\bar{h}^\omega) = 1$ for all ω , using the already proved $\bar{\eta} = \eta$, we have the equality

$$\frac{Q_n^\omega(\bar{h}^{\theta^n\omega})(x)}{\prod_{p=0}^{n-1} \lambda_{\theta^p\omega}} - h^\omega(x) \eta^{\theta^n\omega}(\bar{h}^{\theta^n\omega}) = \bar{h}^\omega(x) - h^\omega(x),$$

so taking $\varphi = \bar{h}^{\theta^n \omega} / \bar{\eta}^{\theta^n \omega} (\bar{h}^{\theta^n \omega})$ in (86), and noting that under **(H2)**,

$$\frac{\|\bar{h}^\omega\|}{\bar{\eta}^\omega(\bar{h}^\omega)} \leq \sup_{\omega, x, x'} \frac{\bar{h}^\omega(x)}{\bar{h}^\omega(x')} = \sup_{\omega, x, x'} \frac{Q(\bar{h}^{\theta^{-1}\omega})(x)}{Q(\bar{h}^{\theta^{-1}\omega})(x')} \leq \beta \frac{\epsilon_+}{\epsilon_-},$$

we find $\bar{h}^\omega(x) = h^\omega(x)$. This completes the proof of part 4), and therefore the proposition. $\square$

Proof. (of Proposition 3)

$$\begin{aligned} \mu Q_n^\omega(1) &= \mu Q_0^\omega(1) \prod_{p=1}^n \frac{\mu Q_p^\omega(1)}{\mu Q_{p-1}^\omega(1)} \\ &= 1 \cdot \prod_{p=1}^n \frac{\mu Q_{p-1}^\omega Q^{\theta^{p-1}\omega}(1)}{\mu Q_{p-1}^\omega(1)} \\ &= \prod_{p=0}^{n-1} \Phi_p^\omega(\mu) Q^{\theta^p \omega}(1) = \prod_{p=0}^{n-1} \Phi_p^\omega(\mu) (G^{\theta^p \omega}), \end{aligned}$$

and so

$$\begin{aligned} \frac{1}{n} \log \mu Q_n^\omega(1) &= \frac{1}{n} \sum_{p=0}^{n-1} \log \lambda_{\theta^p \omega} \\ &+ \frac{1}{n} \sum_{p=0}^{n-1} \left[\log \left[\Phi_p^\omega(\mu) (G^{\theta^p \omega}) \right] - \log \lambda_{\theta^p \omega} \right]. \end{aligned} \quad (87)$$

Now the λ part of (28) ensures that $\mathbb{E}[\log \lambda] < \infty$, so by the ergodic theorem, the first term on the right of (87) converges:

$$\frac{1}{n} \sum_{p=0}^{n-1} \log \lambda_{\theta^p \omega} \rightarrow \mathbb{E}[\log \lambda], \quad \mathbb{P} - a.s.$$

For the other term, fix $\omega \in \Omega$. Then by (28), for all $\epsilon > 0$, there exists $n_0(\epsilon)$ such that for any $\omega' \in \Omega$ and $n > n_0(\epsilon)$,

$$\left| \left[\Phi_n^{\theta^{-n}\omega'}(\mu) (G^{\omega'}) \right] - \lambda_\omega \right| < \epsilon,$$

and then choosing $\omega' = \theta^n \omega$,

$$\left| \left[\Phi_n^\omega(\mu) (G^{\theta^n \omega}) \right] - \lambda_{\theta^n \omega} \right| < \epsilon.$$

Therefore

$$\log \left[\Phi_n^\omega(\mu) (G^{\theta^n \omega}) \right] - \log \lambda_{\theta^n \omega} \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

and so the second term on the right of (87) converges to zero by Cesaro averaging. The proof is complete upon recalling from Proposition 1 that $n^{-1} \log \mu Q_n^\omega(1) \rightarrow \Lambda$, $\mathbb{P} - a.s.$, and noting that by Proposition 2, $Q^\omega(h^{\theta \omega}) = \lambda_\omega h^\omega$. $\square$

Proofs for section 2.4

Throughout this section we assume $N \geq 1$ is fixed arbitrarily.

For $\tilde{\mathbf{M}} \in \mathbb{M}$, define

$$\tilde{\mathbf{R}}(\omega, x, dx') := \mathbf{G}(\omega, x)^2 \phi^\omega(x, x')^2 \tilde{\mathbf{M}}(\omega, x, dx'). \quad (88)$$

The proofs of Proposition 4 and Theorem 1 involve a generalized eigen-value and eigen-function for $\tilde{\mathbf{R}}$, and our first objective is to verify that such quantities exist. In order to do so, we now check that when **(H2)** holds, $\tilde{\mathbf{R}}$ satisfies a regularity condition of a similar form. Define

$$\mathbf{J}(\omega, x) := \int_{\chi^N} \mathbf{G}(\omega, x)^2 \phi^\omega(x, x')^2 \tilde{\mathbf{M}}(\omega, x, dx')$$

and

$$\mathbf{L}(\omega, x, dx') = \frac{\mathbf{G}(\omega, x)^2 \phi^\omega(x, x')^2 \widetilde{\mathbf{M}}(\omega, x, dx')}{\mathbf{J}(\omega, x)},$$

so clearly

$$\widetilde{\mathbf{R}}(\omega, x, dx') = \mathbf{J}(\omega, x) \mathbf{L}(\omega, x, dx').$$

Lemma 6. Assume (H2), let $\widetilde{\mathbf{M}}$ be any member of $\mathbb{M}$ and let $\widetilde{\nu}$ be the accompanying measure in (38). Then there exist constants $\alpha \in [1, \infty)$, $(\delta_-, \delta_+) \in (0, \infty)^2$ and $\mu \in \mathcal{P}(\mathbf{X}^N)$, such

$$\frac{\mathbf{J}(\omega, x)}{\mathbf{J}(\omega', x')} \leq \alpha, \quad \forall (\omega, \omega', x, x') \in \Omega^2 \times \mathbf{X}^{2N}, \quad (89)$$

$$\mu(dx) \propto \left[\frac{d\nu^{\otimes N}}{d\widetilde{\nu}}(x) \right]^2 \widetilde{\nu}(dx) \quad (90)$$

and

$$\delta_- \mu(\cdot) \leq \mathbf{L}(\omega, x, \cdot) \leq \delta_+ \mu(\cdot), \quad \forall (\omega, x) \in \Omega \times \mathbf{X}^N. \quad (91)$$

Proof. For any $A \in \mathcal{X}^{\otimes N}$,

$$\begin{aligned} & \int_A \mathbf{G}^\omega(x)^2 \phi^\omega(x, x')^2 \widetilde{\mathbf{M}}^\omega(x, dx') \\ &= \mathbf{G}^\omega(x)^2 \int_A \left[\frac{d\mathbf{M}^\omega(x, \cdot)}{d\nu^{\otimes N}}(x') \frac{d\nu^{\otimes N}}{d\widetilde{\nu}}(x') \frac{d\widetilde{\nu}}{d\widetilde{\mathbf{M}}^\omega(x, \cdot)}(x') \right]^2 \widetilde{\mathbf{M}}^\omega(x, dx') \\ &\leq \sup_{(\omega', z) \in \Omega \times \mathbf{X}^N} \mathbf{G}^{\omega'}(z)^2 \frac{\beta^2 \epsilon_+^{2N} \widetilde{\epsilon}_+}{\widetilde{\epsilon}_-^2} \int_A \left[\frac{d\nu^{\otimes N}}{d\widetilde{\nu}}(x') \right]^2 \widetilde{\nu}(dx') < \infty, \end{aligned} \quad (92)$$

where Lemma 1, (H2) and the definition of $\mathbb{M}$ have been used. By a similar argument,

$$\begin{aligned} & \int_A \mathbf{G}^\omega(x)^2 \phi^\omega(x, x')^2 \widetilde{\mathbf{M}}^\omega(x, dx') \\ &\geq \inf_{(\omega', z) \in \Omega \times \mathbf{X}^N} \mathbf{G}^{\omega'}(z)^2 \frac{\epsilon_-^{2N} \widetilde{\epsilon}_-}{\beta^2 \widetilde{\epsilon}_+^2} \int_A \left[\frac{d\nu^{\otimes N}}{d\widetilde{\nu}}(x') \right]^2 \widetilde{\nu}(dx'). \end{aligned} \quad (93)$$

Taking $A = \mathbf{X}^N$ in (92) and (93), the bound of (89) holds with

$$\alpha = \beta^6 \frac{\epsilon_+^{2N} \widetilde{\epsilon}_+^3}{\epsilon_-^{2N} \widetilde{\epsilon}_-^3}.$$

The bound of (91) holds with

$$\begin{aligned} \delta_- &= \inf_{(\omega', z) \in \Omega \times \mathbf{X}^N} \mathbf{G}^{\omega'}(z)^2 \frac{\epsilon_-^{2N} \widetilde{\epsilon}_-}{\beta^2 \widetilde{\epsilon}_+^2} \left[\int_{\mathbf{X}} \left[\frac{d\nu^{\otimes N}}{d\widetilde{\nu}}(x') \right]^2 \widetilde{\nu}(dx') \right]^{-1}, \\ \delta_+ &= \sup_{(\omega', z) \in \Omega \times \mathbf{X}^N} \mathbf{G}^{\omega'}(z)^2 \frac{\beta^2 \epsilon_+^{2N} \widetilde{\epsilon}_+}{\widetilde{\epsilon}_-^2} \left[\int_{\mathbf{X}} \left[\frac{d\nu^{\otimes N}}{d\widetilde{\nu}}(x') \right]^2 \widetilde{\nu}(dx') \right]^{-1}. \end{aligned}$$

□

Remark 9. Having established the regularity properties (89)-(91), we notice that $\widetilde{\mathbf{R}}$ will have properties which are exactly similar to those properties of Q established in Propositions 1, 2 and 3, which we shall now summarize (we will not write a proof explicitly, since the arguments follow precisely the same programme as in the proofs of the afore-mentioned Propositions).

For any $N \geq 1$ and $\widetilde{\mathbf{M}} \in \mathbb{M}$,

- there exists a constant $\Xi_N \in (-\infty, \infty)$ such that for any $\mu \in \mathcal{P}(\mathbf{X}^N)$,

$$\frac{1}{n} \log \mu \widetilde{\mathbf{R}}_n^\omega(1) \rightarrow \Xi_N, \quad \text{as } n \rightarrow \infty, \quad \text{for } \mathbb{P} - a.a. \omega. \quad (94)$$

- there exists a random variable $\xi : \Omega \rightarrow \mathbb{R}_+$ (depending on N) and a function $\ell : \Omega \times \mathbf{X}^N \rightarrow \mathbb{R}_+$ which is measurable w.r.t. $\mathcal{F} \otimes \mathcal{X}^{\otimes N}$ and such that

$$\sup_{\omega, \omega'} \frac{\xi_\omega}{\xi_{\omega'}} < \infty, \quad \sup_{\omega, \omega', x, x'} \frac{\ell(\omega, x)}{\ell(\omega, x')} < \infty, \quad (95)$$

and

$$\widetilde{\mathbf{R}}^\omega(\ell^{\theta\omega}) = \xi_\omega \ell^\omega \quad (96)$$

- for any $x \in \mathbf{X}^N$,

$$\Xi_N = \mathbb{E}[\log \xi] = \int_{\Omega} \frac{\widetilde{\mathbf{R}}^\omega(\ell^{\theta\omega})(x)}{\ell^\omega(x)} \mathbb{P}(d\omega). \quad (97)$$

Note that in the above displays, the dependence of various quantities on $\widetilde{\mathbf{M}}$ is suppressed from the notation.

We can now deal with the proof of Proposition 4 and then a collection of Lemmas which prove Theorem 1.

Proof. (of Proposition 4) By Proposition 1, for any $\mu_0 \in \mathcal{P}(\mathbf{X})$,

$$\frac{2}{n} \log \mu_0 Q_n^\omega(1) \rightarrow 2\Lambda, \quad \text{for } \mathbb{P} - a.a. \omega,$$

and by (94),

$$\frac{1}{n} \log \mu_0^{\otimes N} \widetilde{\mathbf{R}}_n^\omega(1) \rightarrow \Xi_N, \quad \text{for } \mathbb{P} - a.a. \omega. \quad (98)$$

Then as

$$\widetilde{\mathcal{V}}_{n,N}^\omega = \frac{\mu_0^{\otimes N} \widetilde{\mathbf{R}}_n^\omega(1)}{[\mu_0 Q_n^\omega(1)]^2},$$

the proof is complete, with $\Upsilon_N(\widetilde{\mathbf{M}}) = \Xi_N - 2\Lambda$. $\square$

The proof of Theorem 1 is now given in three Lemmas. The first can be viewed as generalizing the necessity part of the proof of [Bucklew et al., 1990, Theorem 3] to the case of non-negative kernels driven by an ergodic shift.

Lemma 7. *1) $\Rightarrow$ 2). If $\Upsilon_N(\widetilde{\mathbf{M}}) = 0$, then for $\mathbb{P}$ -almost all $\omega \in \Omega$ there exists $A_\omega \in \mathcal{X}^{\otimes N}$ such that $\nu^{\otimes N}(A_\omega^c) = 0$ and for any $x \in A_\omega$,*

$$\widetilde{\mathbf{M}}^\omega(x, B) = \frac{\int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}, \quad \text{for all } B \in \mathcal{X}^{\otimes N}.$$

Proof. We need to introduce a notational convention before proceeding with the main body of the proof. For any $\varphi : \Omega \times \mathbf{X}^N \rightarrow \mathbb{R}$ a function measurable w.r.t. $\mathcal{F} \otimes \mathcal{X}^{\otimes N}$, let the $\mathbb{P}$ -essential supremum of the collection of functions $\{\varphi(\cdot, x); x \in \mathbf{X}^N\}$ (in the sense of [Doob, 1994, V, 18.]) be say χ , i.e. χ is a random variable on $(\Omega, \mathcal{F}, \mathbb{P})$. In a slight abuse of our ω -section notation we shall write, for any ω in Ω ,

$$\text{ess sup}_x \varphi^\omega(x) := \chi_\omega. \quad (99)$$

Let $\gamma(\omega, x) = \mathbf{h}(\omega, x)^2 / \ell(\omega, x)$, let ρ be the $\mathbb{P}$ -essential supremum of the collection of functions $\{\gamma(\cdot, x) ; x \in \mathbf{X}^N\}$, and then in accordance with the convention (99) we shall write:

$$\operatorname{ess\,sup}_x \frac{\mathbf{h}^\omega(x)^2}{\ell^\omega(x)} := \rho_\omega.$$

From (95), the definition of $\mathbf{h}$ in (95) and (30),

$$\sup_{\omega, \omega', x, x'} \frac{\gamma(\omega, x)}{\gamma(\omega', x')} = \sup_{\omega, \omega', x, x'} \frac{\mathbf{h}^\omega(x)^2}{\ell^\omega(x)} \frac{\ell^{\omega'}(x')}{\mathbf{h}^{\omega'}(x')^2} < \infty \quad (100)$$

and therefore, at least up to a set of $\mathbb{P}$ -measure zero, ρ_ω is uniformly bounded above and below away from zero in ω . This observation, the bound (100) and Lemma 6 will ensure that various expectations appearing below are finite.

We now proceed with the proof. Since by Proposition 2 and (95), λ and ξ are uniformly bounded above and below away from zero in ω , we may write $\Upsilon_N(\widetilde{\mathbf{M}}) = \Xi_N - 2\Lambda = \mathbb{E} \left[\log \frac{\xi}{\lambda^2} \right]$ (where the first equality is as in the above proof of Proposition 4). We are going to prove that the condition $\mathbb{E} \left[\log \frac{\xi}{\lambda^2} \right] = 0$ implies for $\mathbb{P}$ -almost all $\omega \in \Omega$ there exists $A_\omega \in \mathcal{X}^{\otimes N}$ such that $\nu^{\otimes N}(A_\omega^c) = 0$ and for any $x \in A_\omega$,

$$\widetilde{\mathbf{M}}^\omega(x, B) = \frac{\int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}, \quad \text{for all } B \in \mathcal{X}^{\otimes N}. \quad (101)$$

The following holds for $\mathbb{P}$ -almost all ω and any $x \in \mathbf{X}^N$,

$$\begin{aligned} \xi_\omega &= \frac{\widetilde{\mathbf{R}}^\omega(\ell^{\theta\omega})(x)}{\ell^\omega(x)} \\ &\geq \frac{1}{\rho_{\theta\omega} \ell^\omega(x)} \int \mathbf{G}^\omega(x)^2 \phi^\omega(x, x')^2 \widetilde{\mathbf{M}}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \\ &\geq \frac{1}{\rho_{\theta\omega} \ell^\omega(x)} \left[\int \mathbf{G}^\omega(x) \phi^\omega(x, x') \widetilde{\mathbf{M}}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x') \right]^2 \end{aligned} \quad (102)$$

$$\begin{aligned} &= \frac{1}{\rho_{\theta\omega} \ell^\omega(x)} \left[\int \mathbf{Q}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x') \right]^2 \\ &= \frac{\mathbf{h}^\omega(x)^2}{\rho_{\theta\omega} \ell^\omega(x)} \lambda_\omega^2. \end{aligned} \quad (103)$$

where the first equality is just (96), the inequality (102) is due to Jensen's inequality and the equality in (103) holds due to Lemma 2. Therefore the following inequalities hold for $\mathbb{P}$ -almost all ω :

$$\begin{aligned} \xi_\omega &\geq \frac{1}{\rho_{\theta\omega}} \operatorname{ess\,sup}_x \frac{1}{\ell^\omega(x)} \int \mathbf{G}^\omega(x)^2 \phi^\omega(x, x')^2 \widetilde{\mathbf{M}}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \\ &\geq \frac{\lambda_\omega^2}{\rho_{\theta\omega}} \operatorname{ess\,sup}_x \frac{\mathbf{h}^\omega(x)^2}{\ell^\omega(x)}. \end{aligned} \quad (104)$$

Using consecutively the inequalities in (104),

$$\begin{aligned}
\mathbb{E}[\log \xi] &\geq \int_{\Omega} \log \left[\frac{1}{\rho_{\theta\omega}} \operatorname{ess\,sup}_x \frac{1}{\ell^{\omega}(x)} \int \mathbf{G}^{\omega}(x)^2 \phi^{\omega}(x, x')^2 \widetilde{\mathbf{M}}^{\omega}(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \right] \mathbb{P}(d\omega) \\
&= \int_{\Omega} \left[\log \left[\operatorname{ess\,sup}_x \frac{1}{\ell^{\omega}(x)} \int \mathbf{G}^{\omega}(x)^2 \phi^{\omega}(x, x')^2 \widetilde{\mathbf{M}}^{\omega}(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \right] - \log \rho_{\theta\omega} \right] \mathbb{P}(d\omega) \\
&= \int_{\Omega} \log \left[\operatorname{ess\,sup}_x \frac{1}{\ell^{\omega}(x)} \int \mathbf{G}^{\omega}(x)^2 \phi^{\omega}(x, x')^2 \widetilde{\mathbf{M}}^{\omega}(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \right] \mathbb{P}(d\omega) - \mathbb{E}[\log \rho] \quad (105)
\end{aligned}$$

$$\begin{aligned}
&= \int_{\Omega} \log \left[\frac{1}{\rho_{\omega}} \operatorname{ess\,sup}_x \frac{1}{\ell^{\omega}(x)} \int \mathbf{G}^{\omega}(x)^2 \phi^{\omega}(x, x')^2 \widetilde{\mathbf{M}}^{\omega}(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \right] \mathbb{P}(d\omega) \\
&\geq \int_{\Omega} \log \left[\frac{\lambda_{\omega}^2}{\rho_{\omega}} \operatorname{ess\,sup}_x \frac{\mathbf{h}^{\omega}(x)^2}{\ell^{\omega}(x)} \right] \mathbb{P}(d\omega) \quad (106)
\end{aligned}$$

$$= \mathbb{E}[\log \lambda^2] = \mathbb{E}[\log \xi], \quad (107)$$

where (105) holds as θ preserves $\mathbb{P}$; (106) holds due to (104) and because θ preserves $\mathbb{P}$; and the final equality in (107) holds by hypothesis (since we are trying to prove (101)). Thus we conclude

$$\begin{aligned}
\mathbb{E}[\log \xi] &= \int_{\Omega} \log \left[\frac{1}{\rho_{\omega}} \operatorname{ess\,sup}_x \frac{1}{\ell^{\omega}(x)} \int \mathbf{G}^{\omega}(x)^2 \phi^{\omega}(x, x')^2 \widetilde{\mathbf{M}}^{\omega}(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \right] \mathbb{P}(d\omega) \\
&= \int_{\Omega} \log \left[\frac{1}{\rho_{\omega}} \operatorname{ess\,sup}_x \frac{\widetilde{\mathbf{R}}^{\omega} \left([\mathbf{h}^{\theta\omega}]^2 \right) (x)}{\ell^{\omega}(x)} \right] \mathbb{P}(d\omega) \quad (108)
\end{aligned}$$

Now for $\epsilon > 0$ and $\omega \in \Omega$, introduce

$$A_{\omega, \epsilon} := \left\{ x : \ell^{\omega}(x) < \frac{(1 + \epsilon)}{\rho_{\omega}} \mathbf{h}^{\omega}(x)^2 \right\}.$$

Then for $\mathbb{P}$ -almost all ω and all $x \in \mathbf{X}^N$,

$$\ell^{\omega}(x) \geq \frac{\mathbf{h}^{\omega}(x)^2}{\rho_{\omega}} + \epsilon \frac{\mathbf{h}^{\omega}(x)^2}{\rho_{\omega}} \mathbb{I}_{A_{\omega, \epsilon}^c}(x).$$

Now by Proposition 2 and the definition of $\mathbf{h}$, we know $\inf_{\omega, x} \mathbf{h}(\omega, x) > 0$, and similarly, by (95), $\inf_{\omega, x} \ell(\omega, x) > 0$. Combined with Lemma 6 this ensures that there is a strictly positive constant k independent of ϵ and ω such that

$$\begin{aligned}
\xi_{\omega} = \frac{\widetilde{\mathbf{R}}^{\omega}(\ell^{\theta\omega})(x)}{\ell^{\omega}(x)} &\geq \frac{\widetilde{\mathbf{R}}^{\omega}([\mathbf{h}^{\theta\omega}]^2)(x)}{\rho_{\theta\omega} \ell^{\omega}(x)} + \frac{\epsilon}{\rho_{\theta\omega} \ell^{\omega}(x)} \int_{A_{\theta\omega, \epsilon}^c} \widetilde{\mathbf{R}}^{\omega}(x, dx') \mathbf{h}^{\theta\omega}(x')^2 \\
&\geq \frac{\widetilde{\mathbf{R}}^{\omega}([\mathbf{h}^{\theta\omega}]^2)(x)}{\rho_{\theta\omega} \ell^{\omega}(x)} + \frac{\epsilon}{\rho_{\theta\omega}} k \mu(A_{\theta\omega, \epsilon}^c), \quad \forall x \in \mathbf{X}^N,
\end{aligned}$$

where μ is as in Lemma 6. Thus we find, once again using the fact that θ preserves $\mathbb{P}$,

$$\mathbb{E}[\log \xi] \geq \int_{\Omega} \log \left[\frac{1}{\rho_{\omega}} \operatorname{ess\,sup}_x \frac{\widetilde{\mathbf{R}}^{\omega}([\mathbf{h}^{\theta\omega}]^2)(x)}{\ell^{\omega}(x)} + \frac{\epsilon}{\rho_{\omega}} k \mu(A_{\theta\omega, \epsilon}^c) \right] \mathbb{P}(d\omega),$$

but we have already proved (108), and since ϵ , k and ρ_{ω} are strictly positive and finite we deduce that

$$\mathbb{P}(\{\omega : \mu(A_{\theta\omega, \epsilon}^c) = 0\}) = 1. \quad (109)$$

Now for any $\omega \in \Omega$, if $\mu(A_{\theta\omega, \epsilon}^c) = 0$, then by (90) $(d\nu^{\otimes N}/d\tilde{\nu})(x)$ is zero for $\tilde{\nu}$ -a.a. $x \in A_{\theta\omega, \epsilon}^c$, and then $\nu^{\otimes N}(A_{\theta\omega, \epsilon}^c) = 0$. Therefore from (109) we find

$$1 = \mathbb{P}(\{\omega : \nu^{\otimes N}(A_{\theta\omega, \epsilon}^c) = 0\}) = \mathbb{P}(\{\omega : \nu^{\otimes N}(A_{\theta\omega, \epsilon}) = 1\}) = \mathbb{P}(\{\omega : \nu^{\otimes N}(A_{\omega, \epsilon}) = 1\}), \quad (110)$$

where the final equality holds since θ preserves $\mathbb{P}$.

From the definition of $A_{\omega,\epsilon}$ it is clear that for any ω ,

$$\bigcap_{m \geq 1} A_{\omega,1/m} = \left\{ x : \ell^\omega(x) = \frac{1}{\rho_\omega} \mathbf{h}^\omega(x)^2 \right\} =: A_\omega^*, \quad (111)$$

and

$$A_{\omega,1/(m+1)} \subseteq A_{\omega,1/m}, \quad \forall m \geq 1. \quad (112)$$

Now suppose that for some $\omega \in \Omega$, $\nu(A_{\omega,1/m}) = 1$ for all $m \geq 1$. Then in light of (111) and (112), continuity of probability under $\nu^{\otimes N}$ dictates $\nu^{\otimes N}(A_\omega^*) = 1$. On the other hand, if $\nu^{\otimes N}(A_\omega^*) = 1$, then it must be that $\nu^{\otimes N}(A_{\omega,1/m}) = 1$ for all $m \geq 1$. Therefore

$$\{\omega : \nu^{\otimes N}(A_\omega^*) = 1\} = \bigcap_{m \geq 1} \{\omega : \nu^{\otimes N}(A_{\omega,1/m}) = 1\}. \quad (113)$$

Evaluating $\nu^{\otimes N}$ on the sets in (112) we find

$$\{\omega : \nu^{\otimes N}(A_{\omega,1/(m+1)}) = 1\} \subseteq \{\omega : \nu^{\otimes N}(A_{\omega,1/m}) = 1\}, \quad \forall m \geq 1. \quad (114)$$

Then by (110), (113), (114) and continuity of probability under $\mathbb{P}$,

$$\mathbb{P}(\{\omega : \nu^{\otimes N}(A_\omega^*) = 1\}) = 1. \quad (115)$$

Now (115) ensures that, for $\mathbb{P}$ -a.a. ω , A_ω^* is non-empty, so it is legitimate to write, for any $x \in A_\omega^*$,

$$\begin{aligned} \frac{\xi_\omega}{\lambda_\omega^2} &= \frac{R^\omega(\ell^{\theta\omega})(x)}{\ell^\omega(x)} \frac{\mathbf{h}^\omega(x)^2}{[\mathbf{Q}^\omega(\mathbf{h}^{\theta\omega})(x)]^2} \\ &= \frac{\rho_\omega}{\rho_{\theta\omega}} \frac{R^\omega([\mathbf{h}^{\theta\omega}]^2)(x)}{[\mathbf{Q}^\omega(\mathbf{h}^{\theta\omega})(x)]^2} \\ &= \frac{\rho_\omega}{\rho_{\theta\omega}} \frac{\mathbf{G}^\omega(x)^2 \int \phi^\omega(x, x')^2 \widetilde{\mathbf{M}}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')^2}{\mathbf{G}^\omega(x)^2 \left(\int \phi^\omega(x, x') \widetilde{\mathbf{M}}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x') \right)^2} \\ &\geq \frac{\rho_\omega}{\rho_{\theta\omega}}, \end{aligned} \quad (116)$$

where the final inequality is due to Jensen's inequality. Therefore we have proved

$$\frac{\xi_\omega}{\lambda_\omega^2} \frac{\rho_{\theta\omega}}{\rho_\omega} \geq 1, \quad \mathbb{P} - a.a. \omega.$$

However, by hypothesis, $\mathbb{E} \left[\log \frac{\xi_\omega}{\lambda_\omega^2} \right] = 0$, and combined with the fact that θ preserves $\mathbb{P}$ we find

$$\mathbb{E} \left[\log \left(\frac{\xi_\omega}{\lambda_\omega^2} \frac{\rho_{\theta\omega}}{\rho_\omega} \right) \right] = \mathbb{E} \left[\log \frac{\xi_\omega}{\lambda_\omega^2} \right] + \mathbb{E} [\log \rho_{\theta\omega}] - \mathbb{E} [\log \rho_\omega] = 0,$$

therefore it must be the case that in fact

$$\frac{\xi_\omega}{\lambda_\omega^2} \frac{\rho_{\theta\omega}}{\rho_\omega} = 1, \quad \mathbb{P} - a.a. \omega. \quad (117)$$

Equation (117) implies equality must hold in the instance of Jensen's inequality (116), i.e. for $\mathbb{P}$ -almost all ω , and all $x \in A_\omega^*$,

$$\phi^\omega(x, x') \mathbf{h}^{\theta\omega}(x') = c^\omega(x), \quad \widetilde{\mathbf{M}}^\omega(x, \cdot) - a.s.,$$

and therefore

$$\widetilde{\mathbf{M}}^\omega(x, B) = \frac{1}{c^\omega(x)} \int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x'), \quad \text{for any } B \in \mathcal{X}^{\otimes N}.$$

Normalization then dictates

$$\widetilde{\mathbf{M}}^\omega(x, B) = \frac{\int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)},$$

and this completes the proof. $\square$

Lemma 8. *2) $\Rightarrow$ 3). If for $\mathbb{P}$ -almost all $\omega \in \Omega$ there exists $A_\omega \in \mathcal{X}^{\otimes N}$ such that $\nu^{\otimes N}(A_\omega^c) = 0$ and for any $x \in A_\omega$,*

$$\widetilde{\mathbf{M}}^\omega(x, B) = \frac{\int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{X^N} \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}, \quad \text{for all } B \in \mathcal{X}^{\otimes N}.$$

then $\sup_n \widetilde{\mathcal{V}}_{n,N}^\omega < \infty$.

Proof. When

$$\widetilde{\mathbf{M}}^\omega(x, dx') = \frac{\mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{X^N} \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}, \quad (118)$$

for all ω and x , we have already mentioned that $\widetilde{\mathbf{M}} \in \mathbb{M}$ and seen in (44) that

$$\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p\omega}(\zeta_p) \phi^{\theta^p\omega}(\zeta_p, \zeta_{p+1}) = \frac{\mathbf{h}^\omega(\zeta_0)}{\mathbf{h}^{\theta^n\omega}(\zeta_n)} \prod_{p=0}^{n-1} \lambda_{\theta^p\omega}.$$

If (118) holds for some ω, x then we notice that $\widetilde{\mathbf{M}}^\omega(x, \cdot)$ and $\mathbf{M}^\omega(x, \cdot)$ are mutually absolutely continuous, because $\mathbf{h}$ is strictly positive. Therefore if (118) holds only for ω in the set of full $\mathbb{P}$ -measure where the hypothesis of the Lemma holds, and only for $x \in A_\omega$, then using the assumption that θ preserves $\mathbb{P}$, we find that $\nu^{\otimes N}$ -null sets have no contribution to the following expectation:

$$\begin{aligned} \mu_0 \widetilde{\mathbf{R}}_n^\omega(1) &= \widetilde{\mathbb{E}}_N^\omega \left[\prod_{p=0}^{n-1} \mathbf{G}^{\theta^p\omega}(\zeta_p)^2 \phi^{\theta^p\omega}(\zeta_p, \zeta_{p+1})^2 \right] \\ &= \left(\prod_{p=0}^{n-1} \lambda_{\theta^p\omega} \right)^2 \widetilde{\mathbb{E}}_N^\omega \left[\frac{\mathbf{h}^\omega(\zeta_0)^2}{\mathbf{h}^{\theta^n\omega}(\zeta_n)^2} \right] \\ &\leq \left(\prod_{p=0}^{n-1} \lambda_{\theta^p\omega} \right)^2 \left(\sup_{(\omega, \omega', x, x')} \frac{h^\omega(x)}{h^{\omega'}(x')} \right)^2. \end{aligned} \quad (119)$$

Furthermore, since $G^\omega(x) = \lambda_\omega h^\omega(x) / M^\omega(h^{\theta\omega})(x)$,

$$\begin{aligned} [\mu_0 Q_n^\omega(1)]^2 &= \mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^p\omega}(X_p) \right]^2 \\ &= \left(\prod_{p=0}^{n-1} \lambda_{\theta^p\omega} \right)^2 \mathbb{E}^\omega \left[\prod_{p=0}^{n-1} \frac{h^{\theta^p\omega}(X_p)}{M^{\theta^p\omega}(h^{\theta^{p+1}\omega})(X_p)} \right]^2 \\ &= \left(\prod_{p=0}^{n-1} \lambda_{\theta^p\omega} \right)^2 \mathbb{E}^\omega \left[\frac{h^\omega(X_0)}{h^{\theta^n\omega}(X_n)} \prod_{p=1}^n \frac{h^{\theta^p\omega}(X_p)}{M^{\theta^{p-1}\omega}(h^{\theta^p\omega})(X_{p-1})} \right]^2 \\ &\geq \left(\prod_{p=0}^{n-1} \lambda_{\theta^p\omega} \right)^2 \left(\inf_{(\omega, \omega', x, x')} \frac{h^\omega(x)}{h^{\omega'}(x')} \right)^2. \end{aligned} \quad (120)$$

Combining (119) with (120), we arrive at

$$\sup_{n \geq 1} \frac{\mu_0 \widetilde{\mathbf{R}}_n^\omega(1)}{[\mu_0 Q_n^\omega(1)]^2} \leq \left(\sup_{(\omega, \omega', x, x')} \frac{h^\omega(x)}{h^{\omega'}(x')} \right)^4 < \infty,$$

where the final inequality is due to Proposition (2). $\square$

Lemma 9. *3) $\Rightarrow$ 1) If for $\mathbb{P}$ -almost all ω , $\sup_n \tilde{\mathcal{V}}_{n,N}^\omega < \infty$, then $\Upsilon_N(\tilde{\mathbf{M}}) = 0$.*

Proof. Obvious. □

Proof. (of Lemma 3)

We first address $1) \Rightarrow 2)$. By Theorem 1, if $\Upsilon_N(\mathbf{M}) = 0$, then by Theorem 1 it must be the case that up to ω and x being in null sets,

$$\mathbf{M}^\omega(x, B) = \frac{\int_B \mathbf{M}^\omega(x, dx') \mathbf{h}^{\theta\omega}(x')}{\int_{\mathcal{X}^N} \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}, \quad \text{for all } B \in \mathcal{X}^{\otimes N}.$$

and therefore using **(H2)**, we find there must exist a random variable, say χ , such that

$$\text{for } \mathbb{P} - \text{a.a. } \omega, \quad \mathbf{h}^\omega(x) = \chi_\omega, \quad \text{for } \nu^{\otimes N} - \text{a.a. } x.$$

Under **(H2)**, the first equation in (31) shows that the probability measure η^ω is equivalent to ν and therefore

$$\text{for } \mathbb{P} - \text{a.a. } \omega, \quad [\eta^\omega]^{\otimes N}(\mathbf{h}^\omega) = \chi_\omega.$$

But by the third equation in (31), we have $[\eta^\omega]^{\otimes N}(\mathbf{h}^\omega) = \eta^\omega(h^\omega) = 1$, and it follows that

$$\text{for } \mathbb{P} - \text{a.a. } \omega, \quad h^\omega(x) = 1, \quad \text{for } \nu - \text{a.a. } x. \quad (121)$$

Now we show $2) \Rightarrow 3)$. Suppose that (121) holds. Then on appropriate sets, $Q^\omega(h^{\theta\omega}) = \lambda_\omega h^\omega$ reduces to

$$G^\omega(x) = \lambda_\omega =: C_\omega.$$

Finally, we show $3) \Rightarrow 1)$. Suppose there exists a random variable $C : \Omega \rightarrow \mathbb{R}_+$ such that

$$\text{for } \mathbb{P} - \text{a.a. } \omega, \quad G^\omega(x) = C_\omega, \quad \text{for } \nu - \text{a.a. } x.$$

Then since by hypothesis $\tilde{\mathbf{M}}^\omega = \mathbf{M}^\omega$ we have $\tilde{\mathcal{V}}_n^{\omega,N} = 1$ for all $n \geq 1$ and for $\mathbb{P} - \text{a.a. } \omega$. Therefore $\Upsilon_N(\mathbf{M}) = \lim_{n \rightarrow \infty} n^{-1} \log \tilde{\mathcal{V}}_n^{\omega,N} = 0$ with $\mathbb{P}$ probability 1. This completes the proof. □

Proofs for section 3

Proof. (of Lemma 4) The proof of (50) is by induction. At rank $n = 0$ the inequality holds trivially since $\hat{\Phi}_0^\omega = \Phi_0^\omega = Id$. Suppose the inequality holds at rank $n - 1$. Then at rank n ,

$$\begin{aligned} |[\Gamma_{N,n}^\omega(\mu) - \Phi_n^\omega(\mu)](\varphi)| &\leq \left| \left[\Gamma_N^{\theta^{n-1}\omega}(\Gamma_{N,n-1}^\omega(\mu)) - \Phi^{\theta^{n-1}\omega}(\Gamma_{N,n-1}^\omega(\mu)) \right](\varphi) \right| \\ &\quad + \left| \left[\Phi^{\theta^{n-1}\omega}(\Gamma_{N,n-1}^\omega(\mu)) - \Phi^{\theta^{n-1}\omega}(\Phi_{n-1}^\omega(\mu)) \right](\varphi) \right| \\ &\leq \frac{1}{N} \left| \left[\hat{\Phi}^{\theta^{n-1}\omega}(\Gamma_{N,n-1}^\omega(\mu)) - \Phi^{\theta^{n-1}\omega}(\Gamma_{N,n-1}^\omega(\mu)) \right](\varphi) \right| \\ &\quad + \left| \frac{\Gamma_{N,n-1}^\omega(\mu) Q^{\theta^{n-1}\omega}(\varphi)}{\Gamma_{N,n-1}^\omega(\mu) Q^{\theta^{n-1}\omega}(1)} \left[1 - \frac{\Gamma_{N,n-1}^\omega(\mu) Q^{\theta^{n-1}\omega}(1)}{\Phi_{n-1}^\omega(\mu) Q^{\theta^{n-1}\omega}(1)} \right] \right| \\ &\quad + \left| \frac{[\Gamma_{N,n-1}^\omega(\mu) - \Phi_{n-1}^\omega(\mu)] Q^{\theta^{n-1}\omega}(\varphi)}{\Phi_{n-1}^\omega(\mu) Q^{\theta^{n-1}\omega}(1)} \right|, \end{aligned}$$

and the result holds by the induction hypothesis and some simple manipulations.

The proof of (51) is also by induction. At time $n = 0$, the random variables $\{\hat{\zeta}_0^i\}_{i=1}^N$ are iid according to μ_0 , so by the Marcinkiewicz-Zygmund (MZ) inequality,

$$\mathbb{E}_N^\omega \left[\left| [\hat{\eta}_0^N - \mu_0](\varphi) \right|^p \right]^{1/p} \leq \|\varphi\| \frac{C_{0,p}^\omega}{\sqrt{N}}.$$

Suppose the inequality holds at rank $n - 1$. Then at rank n ,

$$\mathbb{E}_N^\omega \left[\left| [\hat{\eta}_n^N - \eta_n^\omega](\varphi) \right|^p \right]^{1/p} \leq \mathbb{E}_N^\omega \left[\left| [\hat{\eta}_n^N - \Gamma_N^{\theta^{n-1}\omega}(\hat{\eta}_{n-1}^N)](\varphi) \right|^p \right]^{1/p} \quad (122)$$

$$+ \mathbb{E}_N^\omega \left[\left| [\Gamma_N^{\theta^{n-1}\omega}(\hat{\eta}_{n-1}^N) - \Gamma_N^{\theta^{n-1}\omega}(\eta_{n-1}^\omega)](\varphi) \right|^p \right]^{1/p} \quad (123)$$

$$+ \left| [\eta_n^\omega - \Gamma_N^{\theta^{n-1}\omega}(\eta_{n-1}^\omega)](\varphi) \right|. \quad (124)$$

The term in (122) is dealt with by application of the MZ inequality and the term in (124) is dealt with using (50). For the term in (123),

$$\begin{aligned} \left[\Gamma_N^{\theta^{n-1}\omega}(\hat{\eta}_{n-1}^N) - \Gamma_N^{\theta^{n-1}\omega}(\eta_{n-1}^\omega) \right](\varphi) &= \frac{1}{N} \left[\hat{\Phi}^{\theta^{n-1}\omega}(\hat{\eta}_{n-1}^N) - \hat{\Phi}^{\theta^{n-1}\omega}(\eta_{n-1}^\omega) \right](\varphi) \\ &+ \left(1 - \frac{1}{N} \right) \frac{\hat{\eta}_{n-1}^N Q^{\theta^{n-1}\omega}(\varphi)}{\hat{\eta}_{n-1}^N Q^{\theta^{n-1}\omega}(1)} \left(1 - \frac{\hat{\eta}_{n-1}^N Q^{\theta^{n-1}\omega}(1)}{\eta_{n-1}^\omega Q^{\theta^{n-1}\omega}(1)} \right) \\ &+ \left(1 - \frac{1}{N} \right) [\hat{\eta}_{n-1}^N - \eta_{n-1}^\omega] \frac{Q^{\theta^{n-1}\omega}(\varphi)}{\eta_{n-1}^\omega Q^{\theta^{n-1}\omega}(1)}, \end{aligned}$$

and the result follows by elementary manipulations involving the MZ inequality and the induction hypothesis. $\square$

Proof. (of Lemma 5) For $a \in \mathbb{R}$ let $[a]$ be the integer part of a and let $\{a\} = a - [a]$. Consider the decomposition

$$\sqrt{N} M_n^{\omega, N}(f^N) = \sum_{k=1}^{(n+1)N} U_k^N(f^N)$$

where for any $1 \leq k \leq (n+1)N$ and (p, i) satisfying $k = pN + i$,

$$U_k^N(f^N) = \begin{cases} \frac{1}{\sqrt{N}} \left[f_p^N(\hat{\zeta}_p^1) - \hat{\Phi}^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N)(f_p^N) \right], & k - pN = 1, \\ \frac{1}{\sqrt{N}} \left[f_p^N(\hat{\zeta}_p^i) - \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N)(f_p^N) \right], & k - pN = i > 1. \end{cases}$$

We will establish distributional convergence of the process

$$X_t^N(f^N) := \sum_{k=1}^{[Nt]+N} U_k^N(f^N),$$

to a continuous Gaussian martingale by application of [Jacod and Shiryaev, 2002, Theorem 3.33, p. 478]. The dependence on ω of $U_k^N(f^N)$, $X_t^N(f^N)$ and various other quantities is suppressed from the notation in the remainder of the proof.

For fixed N , let $\mathcal{H}_k^N$ be the σ -algebra generated by the random variables $\hat{\zeta}_p^i$ for any (p, j) such that $pN + i \leq k$. Then

$$\sum_{k=1}^{[Nt]+N} \mathbb{E}_N^\omega [U_k^N(f^{N,i}) U_k^N(f^{N,j}) | \mathcal{H}_{k-1}^N] = \mathbb{C}_t^N(f^{N,i}, f^{N,j}),$$

where

$$\mathbb{C}_t^N(f^{N,i}, f^{N,j}) = C_{[t]}^N(f^{N,i}, f^{N,j}) + \delta C_t^N(f^{N,i}, f^{N,j}),$$

$$\begin{aligned}
& C_n^N(f^{N,i}, f^{N,j}) \\
&= \frac{1}{N} \sum_{p=0}^n \widehat{\Phi}^{\theta^{p-1}\omega}(\widehat{\eta}_{p-1}^N) \left[\left(f_p^{N,i} - \widehat{\Phi}^{\theta^{p-1}\omega}(\widehat{\eta}_{p-1}^N)(f_p^{N,i}) \right) \left(f_p^{N,j} - \widehat{\Phi}^{\theta^{p-1}\omega}(\widehat{\eta}_{p-1}^N)(f_p^{N,j}) \right) \right] \\
&+ \left(1 - \frac{1}{N} \right) \sum_{p=0}^n \Phi^{\theta^{p-1}\omega}(\widehat{\eta}_{p-1}^N) \left[\left(f_p^{N,i} - \Phi^{\theta^{p-1}\omega}(\widehat{\eta}_{p-1}^N)(f_p^{N,i}) \right) \left(f_p^{N,j} - \Phi^{\theta^{p-1}\omega}(\widehat{\eta}_{p-1}^N)(f_p^{N,j}) \right) \right],
\end{aligned}$$

and

$$\begin{aligned}
& \delta C_t^N(f^{N,i}, f^{N,j}) \\
&= \mathbb{I}[[Nt] - N[t] \geq 1] \frac{1}{N} \widehat{\Phi}^{\theta^{[t]}\omega}(\widehat{\eta}_{[t]}^N) \left[\left(f_{[t]+1}^{N,i} - \widehat{\Phi}^{\theta^{[t]}\omega}(\widehat{\eta}_{[t]}^N)(f_{[t]+1}^{N,i}) \right) \left(f_{[t]+1}^{N,j} - \widehat{\Phi}^{\theta^{[t]}\omega}(\widehat{\eta}_{[t]}^N)(f_{[t]+1}^{N,j}) \right) \right] \\
&+ \mathbb{I}[[Nt] - N[t] \geq 2] \frac{([Nt] - N[t] - 1)}{N} \Phi^{\theta^{[t]}\omega}(\widehat{\eta}_{[t]}^N) \left[\left(f_{[t]+1}^{N,i} - \Phi^{\theta^{[t]}\omega}(\widehat{\eta}_{[t]}^N)(f_{[t]+1}^{N,i}) \right) \left(f_{[t]+1}^{N,j} - \Phi^{\theta^{[t]}\omega}(\widehat{\eta}_{[t]}^N)(f_{[t]+1}^{N,j}) \right) \right]
\end{aligned}$$

with the convention $\widehat{\Phi}^{\theta^{-1}\omega}(\widehat{\eta}_{-1}^N) = \Phi^{\theta^{-1}\omega}(\widehat{\eta}_{-1}^N) = \mu_0$.

Now for $n \geq 0$ consider

$$\mathbb{C}_n(f^i, f^j) := \sum_{p=0}^n \eta_p^\omega \left[(f_p^i - \eta_p^\omega(f_p^i))(f_p^j - \eta_p^\omega(f_p^j)) \right].$$

By Corollary 1, for any $n \geq 0$ and $t \in \mathbb{R}_+$, (and using $[Nt]/N \rightarrow t$)

$$\begin{aligned}
C_n^N(f^{N,i}, f^{N,j}) &\longrightarrow \mathbb{C}_n(f^i, f^j) \\
\delta C_t^N(f^{N,i}, f^{N,j}) &\longrightarrow \{t\} \eta_{[t]+1}^\omega \left[\left(f_{[t]+1}^i - \eta_{[t]+1}^\omega(f_{[t]+1}^i) \right) \left(f_{[t]+1}^j - \eta_{[t]+1}^\omega(f_{[t]+1}^j) \right) \right]
\end{aligned}$$

in probability as $N \rightarrow \infty$. Defining

$$\mathbb{C}_t(f^i, f^j) := \mathbb{C}_{[t]}(f^i, f^j) + \{t\} (\mathbb{C}_{[t]+1}(f^i, f^j) - \mathbb{C}_{[t]}(f^i, f^j))$$

we have proved that for any $t \in \mathbb{R}_+$,

$$\sum_{k=1}^{[Nt]+N} \mathbb{E}_N^\omega \left[U_k^N(f^{N,i}) U_k^N(f^{N,j}) | \mathcal{H}_{k-1}^N \right] \longrightarrow \mathbb{C}_t(f^i, f^j).$$

in probability as $N \rightarrow \infty$.

We now need to verify the conditional Lindeberg condition is satisfied: we shall prove that for any $t \in \mathbb{R}_+$ and $\epsilon > 0$,

$$\sum_{k=1}^{[Nt]+N} \mathbb{E}_N^\omega \left[|U_k^N(f^N)|^2 \mathbb{I}[|U_k^N(f^N)| > \epsilon] | \mathcal{H}_{k-1}^N \right] \rightarrow 0, \quad \text{in probability as } N \rightarrow \infty. \quad (125)$$

To this end, fix $t \in \mathbb{R}_+$, set $n = [t]+1$ and define $\|\varphi\|_n := \vee_{0 \leq p \leq n, 1 \leq q \leq m} \|\varphi_{p,q}\|$. Now for any $1 \leq k \leq [Nt]+N$ setting $p = [(k-1)/N]$, we have by hypothesis of the lemma that $\beta_{p,q}^N$ is measurable w.r.t. $\mathcal{H}_{k-1}^N$ for any $q \in \{1, \dots, m\}$, and therefore

$$\begin{aligned}
& \mathbb{E}_N^\omega \left[|U_k^N(f^N)|^2 \mathbb{I}[|U_k^N(f^N)| > \epsilon] | \mathcal{H}_{k-1}^N \right] \\
& \leq \frac{1}{N} 4 \left(\sum_{q=1}^m |\beta_{p,q}^N| \|\varphi_{p,q}\| \right)^2 \mathbb{I} \left[2 \sum_{q=1}^m |\beta_{p,q}^N| \|\varphi_{p,q}\| > \epsilon \sqrt{N} \right] \\
& \leq \frac{1}{N} 4 \|\varphi\|_n^2 \left(\sum_{q=1}^m |\beta_{p,q}^N| \right)^2 \mathbb{I} \left[2 \|\varphi\|_n \sum_{q=1}^m |\beta_{p,q}^N| > \epsilon \sqrt{N} \right] \\
& \leq \frac{8}{\epsilon N^{3/2}} \|\varphi\|_n^3 \left(\sum_{q=1}^m |\beta_{p,q}^N| \right)^3,
\end{aligned}$$

so in turn,

$$\sum_{k=1}^{[Nt]+N} \mathbb{E}_N^\omega \left[|U_k^N(f^N)|^2 \mathbb{I}[|U_k^N(f^N)| > \epsilon] | \mathcal{H}_{k-1}^N \right] \leq \frac{8}{\epsilon \sqrt{N}} \left(\|\varphi\|_n \right)^3 \sum_{p=0}^n \left(\sum_{q=1}^m |\beta_{p,q}^N| \right)^3. \quad (126)$$

The right hand side of (126) converges to zero in probability as $N \rightarrow \infty$ due to (54) and the continuous mapping theorem. This establishes (125), as required.

Therefore $\{X_t^N(f^N); t \in \mathbb{R}_+\}$ converges in law to a continuous Gaussian martingale $\{X_t(f); t \in \mathbb{R}_+\}$ such that for any $t \in \mathbb{R}_+$ and $1 \leq i, j \leq d$,

$$\langle X(f^i), X(f^j) \rangle_t = \mathbb{C}_t(f^i, f^j).$$

The proof is complete upon noting $X_t^N(f^N) = \sqrt{N} M_{[t]}^{\omega, N}(f^N)$. $\square$

Proof. (of Theorem 2) We first address convergence of the unnormalized measures. Consider the decomposition

$$\begin{aligned}
\hat{\gamma}_n^{\omega, N}(\varphi) - \gamma_n^\omega(\varphi) &= \sum_{p=0}^n \hat{\gamma}_p^{\omega, N}(1) \left[\hat{\eta}_p^N - (\Delta_p^{\omega, N})^{-1} \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] Q_{n-p}^{\theta^p\omega}(\varphi) \\
&= \sum_{p=0}^n \hat{\gamma}_p^{\omega, N}(1) \left[\hat{\eta}_p^N - \frac{\hat{\eta}_p^N(\psi^{\theta^p\omega})}{\Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N)(\psi^{\theta^p\omega})} \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] Q_{n-p}^{\theta^p\omega}(\varphi) \\
&= \sum_{p=0}^n \hat{\gamma}_p^{\omega, N}(1) \left[\hat{\eta}_p^N - \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] (f_{p,n}^{\omega, N}) \\
&= \sum_{p=0}^n \hat{\gamma}_p^{\omega, N}(1) \left[\hat{\eta}_p^N - \Gamma_N^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] (f_{p,n}^{\omega, N}) \\
&\quad + \mathcal{R}_n^{\omega, N}
\end{aligned} \quad (127)$$

with the conventions that $\Delta_p^{\omega, N} = 1$, $\Phi^{\theta^{p-1}\omega}(\hat{\eta}_{-1}^N) = \Gamma_N^{\theta^{p-1}\omega}(\hat{\eta}_{-1}^N) = \mu_0$, and where

$$f_{p,n}^{\omega, N} := Q_{n-p}^{\theta^p\omega}(\varphi) - \frac{\psi^{\theta^p\omega}}{\Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N)(\psi^{\theta^p\omega})} \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) Q_{n-p}^{\theta^p\omega}(\varphi)$$

and

$$\begin{aligned}
\mathcal{R}_n^{\omega, N} &:= \sum_{p=0}^n \hat{\gamma}_p^{\omega, N}(1) \left[\Gamma_N^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) - \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] (f_{p,n}^{\omega, N}) \\
&= \frac{1}{N} \sum_{p=0}^n \hat{\gamma}_p^{\omega, N}(1) \left[\hat{\Phi}^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) - \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] (f_{p,n}^{\omega, N}) \\
&= \frac{1}{N} \sum_{p=0}^n \hat{\gamma}_p^{\omega, N}(1) \hat{\Phi}^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) (f_{p,n}^{\omega, N}).
\end{aligned}$$

Corollary 1 and the continuous mapping theorem ensure that $\hat{\gamma}_n^{\omega, N}(1) \rightarrow \gamma_n^\omega(1)$ and $\sqrt{N}\mathcal{R}_n^{\omega, N} \rightarrow 0$ in probability as $N \rightarrow \infty$. Then by Slutsky's lemma, together with Lemma (5) applied to 127, the proof of (55) is complete.

Now for the normalized measures. Consider the decomposition:

$$\begin{aligned}
\hat{\eta}_n^N(\varphi - \eta_n^\omega(\varphi)) &= \hat{\eta}_0^N \bar{Q}_n^\omega(\bar{\varphi}) + \sum_{p=1}^n \hat{\eta}_p^N \bar{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) - \hat{\eta}_{p-1}^N \bar{Q}_{n-p+1}^{\theta^{p-1}\omega}(\bar{\varphi}) \\
&= \mathcal{M}_n^{N, \omega}(\bar{\varphi}) + \mathcal{R}_n^{N, \omega}(\bar{\varphi}),
\end{aligned}$$

where $\bar{\varphi} = \varphi - \eta_n^\omega(\varphi)$ and where

$$\begin{aligned}
\mathcal{M}_n^{N, \omega}(\varphi) &= \sum_{p=0}^n \left[\hat{\eta}_p^N - \Gamma_N^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] \bar{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) \\
\mathcal{R}_n^{N, \omega}(\varphi) &= \sum_{p=1}^n \Gamma_N^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \bar{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) - \hat{\eta}_{p-1}^N \bar{Q}_{n-p+1}^{\theta^{p-1}\omega}(\bar{\varphi}) \\
&= \sum_{p=1}^n \frac{1}{N} \left[\hat{\Phi}^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) - \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \right] \bar{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) \\
&\quad + \left(1 - \frac{\hat{\eta}_{p-1}^N(G^{\theta^{p-1}\omega})}{\eta_{p-1}^\omega(G^{\theta^{p-1}\omega})} \right) \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \bar{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi})
\end{aligned}$$

with the convention $\Gamma_N^{\theta^{-1}\omega}(\hat{\eta}_{-1}^N) = \mu_0$. Lemma 5 provides the desired convergence in distribution of $\mathcal{M}_n^{N, \omega}(\varphi)$, so by Slutsky's lemma, it remains only to prove that $\sqrt{N}\mathcal{R}_n^{N, \omega}(\varphi) \rightarrow 0$ in probability. To this end, notice that

$$\begin{aligned}
&\mathbb{E}_N^\omega [|\mathcal{R}_n^{N, \omega}(\varphi)|] \\
&\leq \sum_{p=0}^n \frac{2}{N} \left\| \bar{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) \right\| \\
&\quad + \sum_{p=0}^n \frac{1}{\eta_{p-1}^\omega(G^{\theta^{p-1}\omega})} \mathbb{E}_N^\omega \left[\left| \eta_{p-1}^\omega(G^{\theta^{p-1}\omega}) - \hat{\eta}_{p-1}^N(G^{\theta^{p-1}\omega}) \right|^2 \right]^{1/2} \mathbb{E}_N^\omega \left[\left| \Phi^{\theta^{p-1}\omega}(\hat{\eta}_{p-1}^N) \bar{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) \right|^2 \right]^{1/2}
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{E}_N^\omega \left[\left| \Phi^{\theta^{p-1}\omega} (\hat{\eta}_{p-1}^N) \overline{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) - \eta_p^\omega \overline{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi}) \right|^2 \right]^{1/2} \\
& \leq \mathbb{E}_N^\omega \left[\left| \frac{\hat{\eta}_{p-1}^N \overline{Q}_{n-p+1}^{\theta^{p-1}\omega}(\bar{\varphi})}{\hat{\eta}_{p-1}^N (G^{\theta^{p-1}\omega})} \right| \left| 1 - \frac{\hat{\eta}_{p-1}^N (G^{\theta^{p-1}\omega})}{\eta_{p-1}^\omega (G^{\theta^{p-1}\omega})} \right|^2 \right]^{1/2} \\
& \quad + \frac{1}{\eta_{p-1}^\omega (G^{\theta^{p-1}\omega})} \mathbb{E}_N^\omega \left[\left| \hat{\eta}_{p-1}^N \overline{Q}_{n-p+1}^{\theta^{p-1}\omega}(\bar{\varphi}) - \eta_{p-1}^\omega \overline{Q}_{n-p+1}^{\theta^{p-1}\omega}(\bar{\varphi}) \right|^2 \right]^{1/2} \\
& \leq \frac{\|\overline{Q}_{n-p}^{\theta^p\omega}(\bar{\varphi})\|}{\eta_{p-1}^\omega (G^{\theta^{p-1}\omega})} \mathbb{E}_N^\omega \left[\left| \eta_{p-1}^\omega (G^{\theta^{p-1}\omega}) - \hat{\eta}_{p-1}^N (G^{\theta^{p-1}\omega}) \right|^2 \right]^{1/2} \\
& \quad + \frac{1}{\eta_{p-1}^\omega (G^{\theta^{p-1}\omega})} \mathbb{E}_N^\omega \left[\left| \hat{\eta}_{p-1}^N \overline{Q}_{n-p+1}^{\theta^{p-1}\omega}(\bar{\varphi}) - \eta_{p-1}^\omega \overline{Q}_{n-p+1}^{\theta^{p-1}\omega}(\bar{\varphi}) \right|^2 \right]^{1/2}.
\end{aligned}$$

Then by application of Lemma 4 we conclude that there exists a constant $b_\omega(n)$ such that

$$N \mathbb{E}_N^\omega [|\mathcal{R}_n^{N,\omega}(\varphi)|] \leq \|\varphi\| b_\omega(n),$$

which implies, via Markov's inequality, that $\mathcal{R}_n^{N,\omega}(\varphi) \rightarrow 0$ in probability, as required. $\square$

We now turn to the proof of Proposition 5. As per (88), we will work with the non-negative kernel

$$\tilde{\mathbf{R}}(\omega, x, dx') = \mathbf{G}(\omega, x)^2 \phi^\omega(x, x')^2 \tilde{\mathbf{M}}(\omega, x, dx'),$$

In order to prove Proposition 5 we shall introduce the kernel

$$\mathbf{S}(\omega, x, dx') := \frac{1}{[\mathbf{h}^\omega(x)]^2} \tilde{\mathbf{R}}(\omega, x, dx') [\mathbf{h}^{\theta\omega}(x')]^2,$$

which may be regarded as a randomized similarity transform of $\tilde{\mathbf{R}}$. Then writing, in the usual fashion,

$$\mathbf{S}_n^\omega := \mathbf{S}^\omega \dots \mathbf{S}^{\theta^{n-1}\omega}, \quad n \geq 1, \quad (128)$$

it is clear that

$$\mathbf{S}_n^\omega(1)(x) = \frac{1}{[\mathbf{h}^\omega(x)]^2} \tilde{\mathbf{R}}_n^\omega \left([\mathbf{h}^{\theta^n\omega}]^2 \right) (x),$$

and we have

Lemma 10. *Assume (H1), (H2) and*

$$\sup_{\omega, \omega', x, x'} \frac{\psi^\omega(x)}{\psi^{\omega'}(x')} < \infty.$$

Then for any $x \in \mathbb{X}^N$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbf{S}_n^\omega(1)(x) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \tilde{\mathbf{R}}_n^\omega(1)(x), \quad \mathbb{P} - a.s.$$

Proof. Using the bound $\sup_{\omega, \omega', x, x'} h^\omega(x)/h^{\omega'}(x')$ of Proposition 2, we have

$$\begin{aligned}
\frac{1}{n} \log \mathbf{S}_n^\omega(1)(x) &= \frac{1}{n} \log \tilde{\mathbf{R}}_n^\omega \left([\mathbf{h}^{\theta^n\omega}]^2 \right) (x) - \frac{2}{n} \log [\mathbf{h}^\omega(x)]^2 \\
&\leq \frac{1}{n} \log \tilde{\mathbf{R}}_n^\omega(1)(x) + \frac{c}{n},
\end{aligned}$$

for some finite constant c which does not depend on ω or n . This bound, combined with a similar lower one, complete the proof. $\square$

The second component in the proof of Proposition 5 is the following Lemma, whose proof we briefly postpone.

Lemma 11. *Assume (H2),*

$$\sup_{x, x'} \frac{\psi^\omega(x)}{\psi^{\omega'}(x')} < \infty,$$

and fix $\omega \in \Omega$. Then for any $x \in \mathbf{X}^N$ and $N \geq 2$,

$$\left| \frac{\mathbf{S}^\omega(1)(x)}{\lambda_\omega^2} - 1 \right| \leq \frac{1}{N-1} \left[2 \sup_{(z, z') \in \mathbf{X}^2} \frac{\psi^{\theta\omega}(z)}{\psi^{\theta\omega}(z')} - 1 \right] \sup_{z \in \mathbf{X}^N} \left(\frac{\mathbf{M}^\omega(\psi^{\theta\omega})(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(z)} \right) \sup_{(z, z') \in \mathbf{X}^2} \left(\frac{h^{\theta\omega}(z)}{\psi^{\theta\omega}(z)} - \frac{h^{\theta\omega}(z')}{\psi^{\theta\omega}(z')} \right),$$

where λ_ω is as in Proposition 2.

Proof. (of Proposition 5). Let us choose the initial distribution μ_0 to be the eigen-measure η^ω defined in Proposition 2. Iterative application of the equation $\eta^\omega Q^\omega = \lambda_\omega \eta^{\theta\omega}$ shows that

$$\mathbb{E}^\omega \left[\prod_{p=0}^{n-1} G^{\theta^{p-1}\omega}(X_p) \right] = \prod_{p=0}^{n-1} \lambda_{\theta^p\omega},$$

and then by Lemma 10, Proposition 4, (128) and the property that θ preserves $\mathbb{P}$, we have

$$\Upsilon_N(\widetilde{\mathbf{M}}) =_{\mathbb{P}\text{-a.s.}} \lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbf{S}_n^\omega(1)(x) - \frac{2}{n} \sum_{p=0}^{n-1} \log \lambda_{\theta^p\omega} \leq \log \left[\mathbb{P} - \text{ess sup}_\omega \left(\sup_{x \in \mathbf{X}} \frac{\mathbf{S}^\omega(1)(x)}{\lambda_\omega} \right) \right].$$

Applying Lemma 11, and noticing $\sup_{z \in \mathbf{X}^N} \mathbf{M}^\omega(\psi^{\theta\omega})(z)/\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(z) \leq \sup_{x \in \mathbf{X}} \psi^{\theta\omega}(x)/h^{\theta\omega}(x)$, the proof is complete. $\square$

Proof. (of Lemma 11). At various places in the proof we shall write, for some suitable function φ , $\text{osc}(\varphi) := \sup_{x, y} |\varphi(x) - \varphi(y)|$.

The starting point is the expression:

$$\begin{aligned} \frac{\mathbf{S}^\omega(1)(x)}{\lambda_\omega^2} - 1 &= \frac{\mathbf{G}^\omega(x)^2}{\lambda_\omega^2 \mathbf{h}^\omega(x)^2} \mathbf{M}^\omega(\psi^{\theta\omega})(x)^2 \int_{\mathbf{X}^N} \widetilde{\mathbf{M}}^\omega(x, dz) \frac{\mathbf{h}^{\theta\omega}(z)^2}{\psi^{\theta\omega}(z)^2} - 1 \\ &= \frac{\mathbf{M}^\omega(\psi^{\theta\omega})(x) \mathbf{G}^\omega(x)^2}{\mathbf{h}^\omega(x)^2 \lambda_\omega^2} \int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dz) \frac{\mathbf{h}^{\theta\omega}(z)^2}{\psi^{\theta\omega}(z)} - 1 \\ &= \frac{\mathbf{M}^\omega(\psi^{\theta\omega})(x)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)^2} \int_{\mathbf{X}^N} \mathbf{M}^\omega(x, dz) \frac{\mathbf{h}^{\theta\omega}(z)^2}{\psi^{\theta\omega}(z)} - 1. \\ &= \int_{\mathbf{X}^N} \frac{\mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \left[\frac{\mathbf{M}^\omega(\psi^{\theta\omega})(x) \mathbf{h}^{\theta\omega}(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x) \psi^{\theta\omega}(z)} - 1 \right] \\ &= \int_{\mathbf{X}^N} \widetilde{\mathbf{M}}^\omega(x, dz) \left[\frac{d\widetilde{\mathbf{M}}_{\text{opt}}^\omega(x, \cdot)}{d\widetilde{\mathbf{M}}^\omega(x, \cdot)}(z) - 1 \right]^2. \end{aligned}$$

where $\mathbf{M}^\omega(\mathbf{h}^{\theta\omega}) = \lambda_\omega \mathbf{h}^\omega / \mathbf{G}^\omega$ has been used, and the final equality, included only for purposes of exposition, is valid with $\widetilde{\mathbf{M}}_{\text{opt}}^\omega(x, dz) := \mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z) / \mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)$.

The main strategy of the proof is to introduce, for each $\omega \in \Omega$, two judiciously chosen Markov kernels, $K^\omega : \mathbf{X}^N \times \mathcal{X}^{\otimes 2} \rightarrow [0, 1]$ and $L^\omega : \mathbf{X}^N \times \mathcal{X}^{\otimes 2} \rightarrow [0, 1]$ such that we can write

$$\begin{aligned} &\int_{\mathbf{X}^N} \frac{\mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \left[\frac{\mathbf{M}^\omega(\psi^{\theta\omega})(x) \mathbf{h}^{\theta\omega}(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x) \psi^{\theta\omega}(z)} - 1 \right] \\ &= \int_{\mathbf{X}^2} [K^\omega(x, dz^{1:2}) - L^\omega(x, dz^{1:2})] f^\omega(x, z^2), \end{aligned} \tag{129}$$

where

$$f^\omega : (x, z) \in \mathbb{X}^N \times \mathbb{X} \mapsto \frac{\mathbf{M}^\omega(\psi^{\theta\omega})(x) h^{\theta\omega}(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x) \psi^{\theta\omega}(z)} - 1 \in \mathbb{R}_+, \quad (130)$$

and such that we can control the magnitude of (129) using an estimate of $\|K^\omega(x, \cdot) - L^\omega(x, \cdot)\|_{tv}$. Now for any ω and x , by definition $\mathbf{M}^\omega(x, \cdot)$ is a symmetric measure. For $1 \leq q \leq N$ and $x \in \mathbb{X}^N$, we shall write $\mathbf{M}_{(q)}^\omega(x, \cdot)$ for the marginal of $\mathbf{M}^\omega(x, \cdot)$ over the first $q \leq N$ coordinates.

The first Markov kernel we introduce is:

$$K^\omega(x, dz^{1:2}) := \sum_{k=1}^N \sum_{l=1}^N \int_{\mathbb{X}^N} \frac{1}{N} \frac{\mathbf{M}^\omega(x, d\mathfrak{z}) h^{\theta\omega}(\mathfrak{z}^k)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \frac{\psi^{\theta\omega}(\mathfrak{z}^l)}{\sum_{j=1}^N \psi^{\theta\omega}(\mathfrak{z}^j)} \delta_{(\mathfrak{z}^k, \mathfrak{z}^l)}(dz^{1:2}),$$

where here, and henceforth, $\mathfrak{z} = (\mathfrak{z}^1, \dots, \mathfrak{z}^N) \in \mathbb{X}^N$. Elementary manipulations then yield

$$\int_{\mathbb{X}^2} K^\omega(x, dz^{1:2}) f^\omega(x, z^2) = \int_{\mathbb{X}^N} \frac{\mathbf{M}^\omega(x, dz) \mathbf{h}^{\theta\omega}(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \left[\frac{\mathbf{M}^\omega(\psi^{\theta\omega})(x) h^{\theta\omega}(z)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x) \psi^{\theta\omega}(z)} - 1 \right],$$

where f^ω is as in (130).

The second Markov kernel is:

$$L^\omega(x, dz^{1:2}) := \frac{\mathbf{M}_{(2)}^\omega(x, dz^{1:2}) h^{\theta\omega}(z^1) \psi^{\theta\omega}(z^2)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x) \mathbf{M}^\omega(\psi^{\theta\omega})(x)},$$

so that then

$$\int_{\mathbb{X}^2} L^\omega(x, dz^{1:2}) f^\omega(x, z^2) = 0.$$

Furthermore, we have:

$$\begin{aligned} & K^\omega(x, dz^{1:2}) \\ &= \frac{1}{N} \sum_{k=1}^N \int_{\mathbb{X}^N} \frac{\mathbf{M}^\omega(x, d\mathfrak{z}) h^{\theta\omega}(\mathfrak{z}^k)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \frac{\psi^{\theta\omega}(\mathfrak{z}^k)}{\sum_{j=1}^N \psi^{\theta\omega}(\mathfrak{z}^j)} \delta_{(\mathfrak{z}^k, \mathfrak{z}^k)}(dz^{1:2}) \\ &+ \left(1 - \frac{1}{N}\right) \frac{\mathbf{M}_{(2)}^\omega(x, dz^{1:2}) h^{\theta\omega}(z^1) \psi^{\theta\omega}(z^2)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \int_{\mathbb{X}^{N-2}} \frac{\mathbf{M}_{(N-2)}^\omega(x, d\mathfrak{z}^{1:N-2})}{N^{-1} \left[\psi^{\theta\omega}(z^1) + \psi^{\theta\omega}(z^2) + \sum_{j=1}^{N-2} \psi^{\theta\omega}(\mathfrak{z}^j) \right]} \\ &\geq \left(1 - \frac{1}{N}\right) \frac{\mathbf{M}_{(2)}^\omega(x, dz^{1:2}) h^{\theta\omega}(z^1) \psi^{\theta\omega}(z^2)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \\ &\quad \cdot \int_{\mathbb{X}^{N-2}} \mathbf{M}_{(N-2)}^\omega(x, d\mathfrak{z}^{1:N-2}) \left[N^{-1} \left(\psi^{\theta\omega}(z^1) + \psi^{\theta\omega}(z^2) + \sum_{j=1}^{N-2} \psi^{\theta\omega}(\mathfrak{z}^j) \right) \right]^{-1}. \end{aligned} \quad (131)$$

Then, noting that $L^\omega(x, \cdot) \ll K^\omega(x, \cdot)$, for $\mathbf{M}_{(2)}^\omega(x, \cdot)$ – almost all $z^{1:2} \in \mathbb{X}^2$,

$$\begin{aligned} & \frac{dL^\omega(x, \cdot)}{dK^\omega(x, \cdot)}(z^{1:2}) \\ &\leq \left(1 - \frac{1}{N}\right)^{-1} \frac{1}{\mathbf{M}^\omega(\psi^{\theta\omega})(x)} \frac{1}{N} \left(\int \mathbf{M}_{(N-2)}^\omega(x, d\mathfrak{z}^{1:N-2}) \left[\psi^{\theta\omega}(z^1) + \psi^{\theta\omega}(z^2) + \sum_{j=1}^{N-2} \psi^{\theta\omega}(\mathfrak{z}^j) \right]^{-1} \right)^{-1} \\ &\leq \left(1 - \frac{1}{N}\right)^{-1} \frac{1}{\mathbf{M}^\omega(\psi^{\theta\omega})(x)} \frac{1}{N} \left[\psi^{\theta\omega}(z^1) + \psi^{\theta\omega}(z^2) + \int \sum_{j=1}^{N-2} \psi^{\theta\omega}(\mathfrak{z}^j) \mathbf{M}_{(N-2)}^\omega(x, d\mathfrak{z}^{1:N-2}) \right] \\ &= 1 + \frac{1}{N-1} \left[\frac{\psi^{\theta\omega}(z^1) + \psi^{\theta\omega}(z^2)}{\mathbf{M}^\omega(\psi^{\theta\omega})(x)} - 1 \right], \end{aligned}$$

where the first inequality uses (131) and the second is due to Jensen's inequality.

Since $L^\omega(x, \cdot) \ll K^\omega(x, \cdot)$ we then have, with $A = \{z^{1:2} \in \mathbf{X}^2 : \frac{dL^\omega(x, \cdot)}{dK^\omega(x, \cdot)}(z^{1:2}) \geq 1\}$,

$$\begin{aligned} \|L^\omega(x, \cdot) - K^\omega(x, \cdot)\|_{tv} &= \int_A \left[\frac{dL^\omega(x, \cdot)}{dK^\omega(x, \cdot)}(z^{1:2}) - 1 \right] K^\omega(x, dz^{1:2}) \\ &\leq \int_A \frac{1}{N-1} \left[\frac{2 \sup_{\mathfrak{z}} \psi^{\theta\omega}(\mathfrak{z})}{\mathbf{M}^\omega(\psi^{\theta\omega})(x)} - 1 \right] K^\omega(x, dz^{1:2}). \\ &\leq \frac{1}{N-1} \left[\frac{2 \sup_{\mathfrak{z}} \psi^{\theta\omega}(\mathfrak{z})}{\mathbf{M}^\omega(\psi^{\theta\omega})(x)} - 1 \right]. \end{aligned} \tag{132}$$

The tv -norm can also be expressed as

$$\|K^\omega(x, \cdot) - L^\omega(x, \cdot)\|_{tv} = \sup_{\{\varphi: \text{osc}(\phi) \leq 1\}} \left| \int [K^\omega(x, dz^{1:2}) - L^\omega(x, dz^{1:2})] \varphi(z^{1:2}) \right|$$

and combining this with (129) and (132) we obtain

$$\begin{aligned} \left| \frac{\mathbf{S}^\omega(1)(x)}{\lambda_\omega^2} - 1 \right| &= \left| \int [K^\omega(x, dz^{1:2}) - L^\omega(x, dz^{1:2})] f^\omega(x, z^2) \right| \\ &\leq \frac{1}{N-1} \left[2 \sup_{z, z' \in \mathbf{X}} \frac{\psi^{\theta\omega}(z)}{\psi^{\theta\omega}(z')} - 1 \right] \text{osc}(f^\omega(x, \cdot)) \\ &\leq \frac{1}{N-1} \left[2 \sup_{z, z' \in \mathbf{X}} \frac{\psi^{\theta\omega}(z)}{\psi^{\theta\omega}(z')} - 1 \right] \sup_x \left(\frac{\mathbf{M}^\omega(\psi^{\theta\omega})(x)}{\mathbf{M}^\omega(\mathbf{h}^{\theta\omega})(x)} \right) \text{osc}\left(\frac{h^{\theta\omega}}{\psi^{\theta\omega}}\right). \end{aligned}$$

□

References

- K.B. Athreya. Change of measures for Markov chains and the llogl theorem for branching processes. *Bernoulli*, pages 323–338, 2000.
- J.A. Bucklew, P. Ney, and J.S. Sadowsky. Monte Carlo simulation and large deviations theory for uniformly recurrent Markov chains. *J. Appl. Probab.*, 20(1):44–59, 1990.
- F. Cérou, P. Del Moral, and A. Guyader. A nonasymptotic variance theorem for unnormalized Feynman Kac particle models. *Ann. Inst. Henri Poincaré Probab. Stat.*, 47(3), 2011.
- N. Chopin. Central limit theorem for sequential Monte Carlo methods and its application to Bayesian inference. *Ann. Statist.*, 32(6):2385–2411, 2004.
- P. Del Moral. *Feynman-Kac Formulae. Genealogical and interacting particle systems with applications*. Probability and its Applications. Springer Verlag, New York, 2004.
- P. Del Moral and A. Guionnet. Central limit theorem for nonlinear filtering and interacting particle systems. *Ann. Appl. Probab.*, 9(2):275–297, 1999.
- P. Del Moral and A. Guionnet. On the stability of interacting processes with applications to filtering and genetic algorithms. *Ann. Inst. Henri Poincaré Probab. Stat.*, 37(2):155–194, 2001.
- P. Del Moral and J. Jacod. Interacting particle filtering with discrete time observations: Asymptotic behaviour in the Gaussian case. In T. Hida, R.L. Karandikar, H. Kunita, B.S. Rajput, S. Watanabe, and J. Xiong, editors, *Stochastics in Finite and Infinite Dimensions: In Honor of Gopinath Kallianpur*. 2001.
- J.L. Doob. *Measure theory*, volume 143 of *Graduate Texts in Mathematics*. Springer, 1994.

- R. Douc and E. Moulines. Limit theorems for weighted samples with applications to sequential Monte Carlo methods. *Ann. Statist.*, 36(5):2344–2376, 2008.
- R. Douc and E. Moulines. Asymptotic properties of the maximum likelihood estimation in misspecified hidden Markov models. *arXiv e-print:1110.0356*, 2011.
- R. Douc, E. Moulines, and J. Olsson. Optimality of the auxiliary particle filter. *Probab. Math. Statist.*, 29(1):1–28, 2009.
- R. Douc, E. Moulines, and J. Olsson. Long-term stability of sequential Monte Carlo methods under verifiable conditions. *arXiv eprint :1203.6898*, 2012.
- A. Doucet, M. Briers, and S. Sénécal. Efficient block sampling strategies for sequential Monte Carlo methods. *J. Comput. Graph. Statist.*, 15(3):693–711, 2006.
- B. Favetto. On the asymptotic variance in the central limit theorem for particle filters. *ESAIM: Probab. Stat.*, 16:151 – 164, September 2012.
- N.J. Gordon, D.J. Salmond, and A.F.M. Smith. Novel approach to nonlinear/non-Gaussian Bayesian state estimation. In *Radar and Signal Processing, IEE Proceedings F*, volume 140, pages 107–113. IET, 1993.
- D.R. Grey. A note on convergence of probability measures. *J. Appl. Probab.*, pages 1055–1058, 2001.
- J. Jacod and A. Shiryaev. *Limit theorems for stochastic processes*. Springer, 2nd edition, 2002.
- A.M. Johansen and A. Doucet. A note on auxiliary particle filters. *Statist. Probab. Lett.*, 78(12):1498–1504, 2008.
- Y. Kifer. Perron-Frobenius theorem, large deviations, and random perturbations in random environments. *Math. Z.*, 222(1):677–698, 1996.
- H.R. Künsch. Recursive Monte Carlo filters: algorithms and theoretical analysis. *Ann. Statist.*, 33(5):1983–2021, 2005.
- B.G. Leroux. Maximum-likelihood estimation for hidden Markov models. *Stochastic Process. Appl.*, 40(1):127–143, 1992.
- M.K. Pitt and N. Shephard. Filtering via simulation: Auxiliary particle filters. *J. Amer. Statist. Assoc.*, pages 590–599, 1999.
- R. van Handel. Uniform time average consistency of Monte Carlo particle filters. *Stochastic Process. Appl.*, 119(11):3835–3861, 2009.
- N. Whiteley. Stability properties of some particle filters. *ArXiv e-print*, 1109.6779, 2011. Submitted.